

# A Study on Morphometry of Spleen in Aborted Foetuses in Relation to Gestational Age

K.Dharmadas, S.Lokanadham and Arun Raj Pusthela \*

Department of Anatomy, Santhiram Medical College and Hospital, Nandyal, Andhrapradesh-518501

DOI:10.21276/sijap.2019.2.7.6

| Received: 20.07.2019 | Accepted: 27.07.2019 | Published: 30.07.2019

\*Corresponding author: Mr.Arun Raj Pusthela

## Abstract

**Background:** Spleen is an essential organ due to its significant role in immunological and haematological functions of the body. Splenic notches on various borders establish the lobulation of the spleen in early life. **Aim:** To study the morphological and morphometric parameters of the spleen in aborted foetuses. **Materials and Methods:** 30 foetal spleen specimens (Male-14; Female-16) of both sexes from 12-40 weeks of gestational age were obtained from the Department of Anatomy, Santhiram Medical College, Nandyal. The morphometric parameters like length, width, thickness and weight of the foetal spleen were measured and recorded. **Results:** The incidence of wedge shape was 23 (76.66%), the pyramidal shape was 6(20%), and leaf-shaped was 1(3.33%). The incidence of the single notch was observed in 19 (63.33%), two notches in 9(30%), and three notches in 2(6.66%) specimens were found in the present study. The parameters like length, width, thickness, and weight of the fetal spleen were increased in relation to the gestational age in the present study. **Conclusion:** The present study acknowledges the anatomical and morphological basis of the spleen may be helpful for the surgeons while doing partial splenectomy in the removal of the affected segment by ligating the segmental branch of the splenic artery.

**Keywords:** Morphometry, notch, shape, spleen.

**Copyright @ 2019:** This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

## INTRODUCTION

Spleen is located in the left hypochondrium and partly in the epigastric region of the abdomen between the fundus of the stomach and the diaphragm [1]. Accessory spleens may be located within the gastrosplenic ligament and at the hilum of the spleen or occasionally, within the greater omentum and rarely left spermatic cord. The stomach and left colic flexure influence the shape of the spleen and three morphological types are desired [2]. The shape of the spleen is crescentic, rhomboid and triangular [3, 4]. The size of spleen varies with age, with the individual and in the same individual under different conditions [5]. Spleen has diaphragmatic and visceral surfaces, superior and anterior borders and inferior and posterior extremities [6]. There is limited research about the sizes of the foetal spleen and infant spleen at 6th week. The length of the spleen is 10mm [5]. The present study was aimed to study the morphological and morphometric parameters of the spleen in aborted foetuses in relation to gestational age.

## MATERIALS AND METHODS

A total of 30 foetal spleen specimens (Male-14; Female-16) obtained from the Department of Anatomy, Santhiram Medical College, Nandyal to study the morphological and morphometric parameters of the foetal spleen in relation to gestational age. The foetuses were of both the sexes and of 12 – 40 weeks of gestational age. The foetuses were preserved by injecting 10%formal solution into the pleural, peritoneal and the cranial cavities. External features of the foetuses and visible anomalies if any were recorded. The morphological parameters like shape, colours were noted, and the morphometric parameters like length, width and thickness of the kidneys by using vernier callipers and recorded. The specimens were categorized into 0-12 weeks, 12-24 weeks, 24-36weeks and >36 weeks of gestational ages. The shapes of the spleen, the number of notches were recorded. Morphometric measurements of fetal spleens were taken with the help of vernier calliper. The weight of the spleens was measured by using a weighing scale. The length of the spleen measured as the distance between the anterior and posterior extremities. The width of the spleen measured as the distance between the superior and

inferior borders. The thickness of the spleen measured as the distance between the superior and inferior borders.

## RESULTS

All the foetal spleen specimens were reddish-brown colour in our study. Out of 30 foetal spleen specimens, 23 specimens were wedge-shaped, and six specimens were pyramidal, and one specimen was leaf-shaped observed in the present study [Table-1]. The incidence of wedge shape was 23(76.66%), the pyramidal shape was 6(20%), and leaf-shaped was 1(3.33%). The incidence of the single notch was observed in 19(63.33%), two notches in 9(30%), and three notches in 2(6.66%) specimens were observed in the present study. The weight of the fetal spleen was  $0.02 \pm 0.01$ gm in 0-12 weeks,  $0.53 \pm 0.048$ gm in 12-24 weeks,  $1.91 \pm 0.07$ gm in 24-36 weeks, and  $4.09 \pm 0.08$ gm in > 36 weeks of gestational ages respectively. The length of fetal spleen was  $5.95 \pm 0.03$ mm, the width was  $2.96 \pm 0.35$  mm, and thickness was  $2.15 \pm 0.62$  mm in 0-12 weeks of gestational age group fetuses. The length of fetal spleen was  $11.73 \pm 0.61$  mm, the width was  $6.79 \pm 1.18$  mm, and thickness was  $5.21 \pm 0.58$  mm in 12-24 weeks of gestational age group fetuses. The length of fetal spleen was  $20.07 \pm 1.24$  mm, the width was  $12.92 \pm 0.62$  mm, and thickness

was  $9.264 \pm 0.66$  mm in 24-36 weeks of gestational age group fetuses. In >36 weeks of gestational age group fetuses, the length of fetal spleen was  $22.41 \pm 0.90$  mm, the width was  $15.65 \pm 0.63$  mm, and thickness was  $10.66 \pm 0.63$  mm noted in the present study[Table-2].



**Fig-1: location of spleen in left hypochondrium in aborted fetus insitu**

**Table-1: Distribution of shape of the fetal spleen in our study**

S.No	Shape of the spleen	Number of specimens
1	Wedge	23
2	Pyramidal	6
4	Leaf shaped	1

**Table-2: Morphometric parameters of the fetal spleen in our study**

Gestational age (Weeks)	Mean±Sd			
	Length(mm)	Width(mm)	Thickness(mm)	Weight of spleen (Gms)
0-12 weeks	$5.95 \pm 0.03$	$2.96 \pm 0.35$	$2.15 \pm 0.62$	$0.02 \pm 0.01$
12-24 weeks	$11.73 \pm 0.61$	$6.79 \pm 1.18$	$5.21 \pm 0.58$	$0.53 \pm 0.048$
24-36 weeks	$20.07 \pm 1.24$	$12.92 \pm 0.62$	$9.264 \pm 0.66$	$1.91 \pm 0.07$
>36 weeks	$22.41 \pm 0.90$	$15.65 \pm 0.63$	$10.66 \pm 0.63$	$4.09 \pm 0.08$

## DISCUSSION

The location of spleen was observed in left hypochondrium of the abdomen in all the fetuses (Figure-1) in our study in agreement with previous literature [6]. Development and morphometry of spleen studied by the use of ultrasonography in the majority of the studies [7,8]. There was a difference in morphometric parameters of spleens between alive and dead subjects [9]. Three faced wedge-shaped fetal spleen in 60% cases; four-faced in 36.5% and two-faced in 3.5% cases [10]. The length, width and thickness of fetal spleen was 1.4 cm, 1.1 cm and 0.8 cm in 12 to 24-week fetuses and 2.4 cm, 1.5 cm and 1.0 cm in 25 to 36-week fetuses[11]. Higher values of dimensions of the fetal spleen as compared to our study probably because of fresh fetuses [12,13]. The size of the spleen increases very rapidly in the embryo from about 6<sup>th</sup> month [14]. The difference in the dimensions of the fetal spleen in our study may be due to the consideration of differential age group and number of fetuses collected for the study. Morphometric parameters of the fetal spleen in our study are increasing in relation to the gestational age of the fetuses was significant in our study, which is in

agreement with previous literature [4]. Limited literature is available on morphometric parameters of the fetal spleen; the research needs to be extended on an increasing number of specimens in relation to each age group may help us in better understanding the morphometry of the spleen for future prospects of the study [15,16].

## CONCLUSION

Our study acknowledges the morphological and morphometric parameters of the spleen may be helpful to the clinicians, radiologist and surgeons while doing partial splenectomy in the removal of the affected segment by ligating the segmental branch of the splenic artery.

## Acknowledgement:

The authors are also grateful to authors, editors and publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed.

## REFERENCES

1. Rosenberg, H. K., Markowitz, R. I., Kolberg, H., Park, C., Hubbard, A., & Bellah, R. D. (1991). Normal splenic size in infants and children: sonographic measurements. *AJR. American journal of roentgenology*, 157(1), 119-121.
2. Sir, A.C., Robert, J.C. (2000). Steel. Essentials of Surgical Practice, Butterworth-Heinemann Limited, 4th edition, 454-75.
3. Prassopoulos, P., Daskalogiannaki, M., Raissaki, M., Hatjidakis, A., & Gourtsoyiannis, N. (1997). Determination of normal splenic volume on computed tomography in relation to age, gender and body habitus. *European radiology*, 7(2), 246-248.
4. Ranganathan, T.S. (2006). A text book of Human Anatomy. S. Chand Group, 281-82.
5. Hostetler, J. R., & Ackerman, G. A. (1969). Lymphopoiesis and lymph node histogenesis in the embryonic and neonatal rabbit. *American Journal of Anatomy*, 124(1), 57-75.
6. Bannister. (1995). Gray's Anatomy 38th edition. Churchill Livingstone, 1437-45.
7. Schmidt, W., Yarkoni, S., Jeanty, P., Grannum, P., & Hobbins, J. C. (1985). Sonographic measurements of the fetal spleen: clinical implications. *Journal of Ultrasound in Medicine*, 4(12), 667-672.
8. Aoki, S., Hata, T., & Kitao, M. (1992). Ultrasonographic assessment of fetal and neonatal spleen. *American journal of perinatology*, 9(05/06), 361-367.
9. Griffith, R. C., & Janney, C. G. (1990). Hematopoietic system: bone marrow and blood, spleen, and lymph nodes. *Anderson's pathology*, 2, 1373-1492.
10. Üngör, B., Malas, M. A., Sulak, O., & Albay, S. (2007). Development of spleen during the fetal period. *Surgical and Radiologic Anatomy*, 29(7), 543-550.
11. Singh, R., & Kumari, G. (2016). Morphological Assessment of Human Foetal Spleen. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 6(52), 19.
12. Soylulu, A., Tanyeli, E., Marur, T., Ertem, A. D., Özkü, K., & Akkn, S. M. (1996). Splenic artery and relation between the tail of the pancreas and spleen in a surgical anatomical view. *Karadeniz Tıp Dergisi*, 9, 103-107.
13. Marecki, B. (1989). The formation of the proportions of the liver, spleen and kidneys in the fetal ontogenesis. *Zeitschrift für Morphologie und Anthropologie*, 117-132.
14. Henry Gary. On the structure and uses of the spleen. JW Parker and Sons. 1860:36.
15. Halsey, C., Rivers, R., Bain, B. J., & Wilkins, B. S. (2004). Unusual splenic sinusoidal iron overload in sickle cell/haemoglobin D-Punjab disease. *Journal of clinical pathology*, 57(5), 539-540.
16. Kuntal, V., Vinayak, G., Kanchan, K. (2018). Morphometric study of human foetal spleen: a study from north-india. *Int J Anat Res*, 6(4.3), 5983-88.