

# Apoptosis Inducing Activity of Proteins Isolated from *Muntingia Calabura* Plant Root on Oral Cancer Cell Line: An *In Vitro* Study

Kumaran C<sup>1</sup>, Dinesha R<sup>2\*</sup>, Santhosh Kumar N<sup>3</sup>

<sup>1</sup>Professor, Department of Pathology, Oxford Medical College & Research Centre, Bangalore, Karnataka, India

<sup>2</sup>Scientific Officer, Adichunchanagiri Institute for Molecular Medicine (AIMM), AIMS- Central Research Laboratory, B.G. Nagara, Mandya, Karnataka, India

<sup>3</sup>Assistant Professor, Department of Biochemistry, Shridevi Institute of Medical Sciences, Tumkur, Karnataka, India

DOI: [10.36348/SJPM.2019.v04i12.001](https://doi.org/10.36348/SJPM.2019.v04i12.001)

| Received: 20.11.2019 | Accepted: 27.11.2019 | Published: 11.12.2019

\*Corresponding author: Dr. Dinesha Ramadas

## Abstract

Now days, research studies have concentrated on complementary (alternate) medicine in treating a large number of infectious and non-infectious diseases, including cancer. The attention of research now shifted towards alternative natural medicine and is being preferred to the toxic effects of the synthetic drugs that are used to treat such diseases. Studies have shown that many natural plant products like amygdalin extracted from apricots and almonds to have shown anticarcinogen effect on many types of cancers including oral cancer. Herein we made an attempt to evaluate the anti-carcinogenic property of *Muntingia Calabura* proteins against Oral cancer cell lines. Oral cancer cell lines (KB cell line) were used in the present study. The proteins were extracted from *Muntingia Calabura* roots and the antiproliferative and cytotoxic activity on KB cell line was evaluated using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide assay. The crude proteins of *Muntingia Calabura* roots showed cytotoxic and anti-proliferative activity on KB cell lines at a maximum efficacy at 100 µg/ml is about 58% and maximum IC50 value is 52 µg/ml. The crude proteins of *Muntingia Calabura* roots effective as an antiproliferative agent, who caused apoptosis in oral cancer cell line.

**Keywords:** *Muntingia Calabura*, plant proteins, oral cancer, squamous cell carcinoma, KB cell lines.

**Copyright © 2019:** This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and sources are credited.

## INTRODUCTION

In oral cancer, the squamous cell carcinoma (SCC) is the very common type of cancer with high rate of mortality and morbidity [1, 2]. In the Indian subcontinent, the oral cancer is occupied the 6<sup>th</sup> position [3]. Various bad habits like chewing of tobacco, betel quid, areca nut [4, 5] and even the infection of human papilloma virus (HPV) also increase the risk factor of Oral Cancer [6, 7]. Though the pharmaceutical industries are developing more potent anticancer drugs with the help of researchers, the conventional treatments such as surgery, radiotherapy and chemotherapy, problems related to these therapies such as side effects, opportunistic infections and the development of drug resistance remained unsolved [8, 10]. Hence, there is an urgency to develop novel treatment ways using plant derivatives which act as effective therapeutic agents and have minimal / no side effects.

Many studies have shown that the different solvent extracts of *Muntingia Calabura* can inhibit /prevent different types of Cancers. Proteins of

*Muntingia Calabura* plant root is one such natural anticancerous plant product, whose anticancer effect on oral cancer has never been reported. Herein the study was designed for the *in vitro* apoptotic effect of *Muntingia Calabura* plant root proteins on SCC cancer cells.

## MATERIALS AND METHODS

### Preparation of Extracts

10g of fresh roots of *Muntingia Calabura* plant were collected from local area, washed thoroughly with distilled water, cut in to small pieces and crushed with 200 ml of double distilled water. Further vortexed for two hours at room temperature, was centrifuged at 10000 rpm at 4°C for 20 minutes. The supernatant collected was subjected to protein precipitation using 55% Ammonium Sulphate. The mixture kept for vortex overnight at 10°C. Further, the mixture was centrifuged at 10,000 rpm using refrigerated centrifuge, the precipitated protein was collected. The collected ammonium sulphate protein precipitate was desalted using molecular cutoff centricons (2kDa). The salt free protein precipitate was stored in deep freezer for further

analysis. Five concentrations of 5, 10, 15, 20 and 25 µg of proteins were prepared.

### Cell Line and Cell Culture

Oral SCC cell line (KB mouth cell line) procured from the National Centre for Cell Sciences, Pune, India for the study. The KB mouth cell line was seeded into a 96 well microtiter plate containing Eagle Minimum Essential Medium supplemented with fetal bovine serum (10%) and penicillin (1%) and streptomycin. With 5% humidity, the cells were maintained in a CO<sub>2</sub> incubator at 37°C for 48 to 72 h.

### Activity of Extracts on Oral Cancer Cell Line

After the incubation (24 hours), the seeded oral SCC cell lines (KB mouth cell line) in 96- well plates at a density of  $5 \times 10^4$  cells/well, were treated with crude proteins of *Muntingia Calabura* plant root of five concentrations viz 5, 10, 15, 20 and 25 µg/mL diluted with dimethyl sulfoxide (DMSO) and incubated for 24 to 48 hours. Based on the mitochondrial dehydrogenase activity in the living cells, the proliferation activity of cell population under different concentrations was determined.


### Cell Viability Assay

The Cell viability assay was done to determine the sublethal concentrations of inhibitory concentration (IC<sub>50</sub>) and proliferative activity of the cells in the presence of crude proteins of *Muntingia Calabura* plant root. After the completion of incubation period, working solution 10 µl of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide was added to each culture well to detect the cell viability. The color was allowed to develop for an additional 4 hours of incubation. Equal volume of DMSO was added to stop this reaction to solubilize the crystals. Absorbance was recorded at 570 nm using Shimadzu UV-Visible spectrophotometer [11].

### STATISTICAL ANALYSIS

The data are given as the mean  $\pm$  standard deviation (SD). The statistical analysis was performed by using one-way analysis of variance (ANOVA), followed by a Dunnett's multiple-comparison test with 95% confidence intervals. A P-value < 0.01 was considered to indicate a statistically significant difference.

## RESULTS



**Fig-1:** KB cell lines as control, and the apoptotic activities of crude proteins of *Muntingia Calabura* plant root against KB cell lines at 50 µg and 100 µg/ml

The crude proteins of *Muntingia Calabura* plant root proteins showed antitumor activity against KB cell line. The viability decreased in a dose-dependent manner. The cell viability decreased as the concentration of the extracts increased. The proteins exhibited their efficacy at 100 µg/mL by killing 58% of cells (Figure-1). The IC<sub>50</sub> values were also calculated, as the crude proteins of *Muntingia Calabura* plant proteins showed a maximum IC<sub>50</sub> at 52 µg/mL. Similar studies have been done using Almond, Apricot extract and Berberine a natural isoquinoline alkaloid isolated from plant genus *Coptis* [12, 13]. The viability decreased in a dose-dependent manner. The crude proteins also showed a promising cytotoxic efficacy.

**Table-1:** The percentage of cell population in different period of time (Cytotoxicity studies)

Percentage	24 Hours	48 Hours
Live	84.11	27.12
Apoptosis	41.04	38.31
Late apoptosis	25.02	50.22

## DISCUSSION

It is reported that, the different parts of *Muntingia Calabura* plant contains significant amount of proteins, tannins, alkaloids, steroids and flavonoids. The methanol extract of *Muntingia Calabura* plant showed antifouling and anti microbial activity in different model systems [14]. Further, it is reported that, the aqueous leaf extract of *Muntingia Calabura* plant at concentrations of 10%, 50% and 100% showed significant antinociceptive, anti-inflammatory and antipyretic activities [15]. It is reported that, the extracts of roots, leaves and fruits of *Muntingia Calabura* plant showing excellent antioxidant activities in different model systems [16]. It is reported that, the flavonoid rich methanolic extract of *Muntingia calabura* plant leaves showed anticarcinogenic activity against the azoxymethane-induced colon cancer in rats involved modulation of the colonic antioxidant system [17]. Furthermore, it is reported that, the antinociceptive activity of MEMC involved activation of the non-selective opioid (particularly the  $\mu$ -,  $\delta$ - and  $\kappa$ -opioid)

and non-opioid (particularly adenosinergic,  $\alpha_2$ -noradrenergic, and  $\beta$ -adrenergic) receptors, modulation of the ATP-sensitive  $K^+$  channel, and inhibition of bradikinin and protein kinase C actions. The discrepancies in MEMC antinociception could be due to the presence of various phytochemicals [18]. With all the above surveys, it was noticed by us is that, no scientific reports or studies are done using proteins isolated from roots of *Muntingia Calabura* plants. The obtained results are very promising and hence further more purification proteins are needed and the studies are continued in this direction.

## CONCLUSION

Based on the results of our study and literature review, the crude proteins extracted from roots of *Muntingia Calabura* plant are showed cytotoxic effect on human oral cancer cell lines. However, purification of these proteins is ongoing to find the protein responsible for this anticancer property.

## REFERENCE

- Nóbrega, T. D., Salomão-Israel-Monteiro-Lourenço Queiroz, E., de Moura Santos, A. D. L., Costa, L., Pereira-Pinto, L., & de Souza, L. B. (2018). Clinicopathological evaluation and survival of patients with squamous cell carcinoma of the tongue. *Medicina oral, patologia oral y cirugia bucal*, 23(5), e579-e587.
- Markopoulos, A. K. (2012). Current aspects on oral squamous cell carcinoma. *The open dentistry journal*, 6, 126-130.
- Coelho, K. R. (2012). Challenges of the oral cancer burden in India. *Journal of cancer epidemiology*, 701932.
- Niaz, K., Maqbool, F., Khan, F., Bahadar, H., Hassan, F. I., & Abdollahi, M. (2017). Smokeless tobacco (paan and gutkha) consumption, prevalence, and contribution to oral cancer. *Epidemiology and health*, 39:e2017009.
- Lee, C. Y., Wu, C. F., Chen, C. M., & Chang, Y. Y. (2018). Qualitative study for betel quid cessation among oral cancer patients. *PloS one*, 13(7), e0199503.
- Kim, S. M. (2016). Human papilloma virus in oral cancer. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*, 42(6), 327-336.
- Candotto, V., Lauritano, D., Nardone, M., Baggi, L., Arcuri, C., Gatto, R., ... & Carinci, F. (2017). HPV infection in the oral cavity: epidemiology, clinical manifestations and relationship with oral cancer. *ORAL & implantology*, 10(3), 209-220.
- Huang, C. Y., Ju, D. T., Chang, C. F., Reddy, P. M., & Velmurugan, B. K. (2017). A review on the effects of current chemotherapy drugs and natural agents in treating non-small cell lung cancer. *Biomedicine*, 7(4):23.
- Casals, E., Gusta, M. F., Cobaleda-Siles, M., Garcia-Sanz, A., & Puentes, V. F. (2017). Cancer resistance to treatment and antiresistance tools offered by multimodal multifunctional nanoparticles. *Cancer nanotechnology*, 8(1), 7.
- Zheng, P. P., Li, J., & Kros, J. M. (2018). Breakthroughs in modern cancer therapy and elusive cardiotoxicity: Critical research- practice gaps, challenges, and insights. *Medicinal research reviews*, 38(1), 325-376.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of immunological methods*, 65(1-2), 55-63.
- Sireesha, D., Reddy, B. S., Reginald, B. A., Samatha, M., & Kamal, F. (2019). Effect of amygdalin on oral cancer cell line: An in vitro study. *Journal of oral and maxillofacial pathology: JOMFP*, 23(1), 104-107.
- Kim, J. S., Oh, D., Yim, M. J., Park, J. J., Kang, K. R., Cho, I. A., ... & Kim, D. K. (2015). Berberine induces FasL-related apoptosis through p38 activation in KB human oral cancer cells. *Oncology reports*, 33(4), 1775-1782.
- Singh, R., Iye, S., Prasad, S., Deshmukh, N., Gupta, U., Zanje, A., ... & Joshi, S. (2017). Phytochemical Analysis of Muntingia calabura Extracts Possessing Anti-Microbial and Anti-Fouling Activities. *International Journal of Pharmacognosy and Phytochemical Research*, 9(6), 826-832.
- Zakaria, Z. A., Hazalin, N. M. N., Zaid, S. M., Ghani, M. A., Hassan, M. H., Gopalan, H. K., & Sulaiman, M. R. (2007). Antinociceptive, anti-inflammatory and antipyretic effects of Muntingia calabura aqueous extract in animal models. *Journal of Natural Medicines*, 61(4), 443-448.
- Nagara, B. G. (2015). Antioxidant Activity: Root, Leaves and Fruits Aqueous Extracts of MuntingiaCalabura. *Journal of Innovations in Pharmaceuticals and Biological Sciences*, 2(4):363-368.
- Md Nasir, N. L., Kamsani, N. E., Mohtarrudin, N., Othman, F., Md. Tohid, S. F., & Zakaria, Z. A. (2017). Anticarcinogenic activity of Muntingia calabura leaves methanol extract against the azoxymethane-induced colon cancer in rats involved modulation of the colonic antioxidant system partly by flavonoids. *Pharmaceutical biology*, 55(1), 2102-2109.
- Zakaria, Z. A., Sani, M. H. M., Cheema, M. S., Kader, A. A., Kek, T. L., & Salleh, M. Z. (2014). Antinociceptive activity of methanolic extract of Muntingia calabura leaves: further elucidation of the possible mechanisms. *BMC complementary and alternative medicine*, 14(1), 63.