

## Research Article

# Comparative Study of Effect of Farming on Anthropological Parameters among Farmers and Non-farmers

Sougata Ganguly<sup>1</sup>, Rajkumar Maiti<sup>2</sup>, Megha Gupta<sup>3</sup>, Prithviraj Karak<sup>2</sup>

<sup>1</sup>Student, Centre for Applied Genetics, Bangalore University, Janna Bharathi, Bangalore-560056

<sup>2</sup>Assistant Professor, Department of Physiology, Bankura Christian College, P.O. + Dist- Bankura, Pin-722101

<sup>3</sup>Student, Department of Biotechnology, Indian Academy Degree College, Hennur Cross, Kalyan Nagar, Bangalore-560043

### \*Corresponding Author:

Prithviraj Karak

Email: [drpkarak@gmail.com](mailto:drpkarak@gmail.com)

**Abstract:** In most countries, agriculture is recognized as one of the most hazardous industries. Investigating the health status of agricultural workers is a challenging goal. The aim of this study was to examine the impact of physical activity and exercise on both farmers' and non-farmers health status on farm production at selected villages in Bankura district, West Bengal. In this cross-sectional study, certain anthropological parameters were compared between farmers and non-farmers living in the same rural area. Farmers and non-farmers, matched per age and sex, were selected randomly in the agricultural areas of different parts of Bankura. According to personal statements, farmers suffered from several musculoskeletal disorders and joint pains such as waist region, low back pain. BMI, HR, BP were found to be lower among farmers. Whereas their PFI scores and recovery duration were higher than non-farmers. This study also shows though farmers have more musculoskeletal disorders than normal sedentary workers, farmers are comparatively more fit than non-farmers. According to personal statements, farmers suffered from hypertension, cardiovascular and orthopaedic problems in higher frequency.

**Keywords:** Farming, health status, anthropological, musculoskeletal disorder, hypertension

## INTRODUCTION

Agriculture plays a pivotal role in the Indian economy. Although its contribution to gross domestic product (GDP) is now around one sixth, it provides employment to 56% of the Indian workforce. Majority of Indian population live in rural areas, mainly depending on agriculture for their livelihood, and carry out more physical activities when compared to urban population who are accustomed to sedentary life style. Healthy body is necessary for increasing the working capacity and maintaining physical fitness of any individual to perform his daily tasks vigorously and alertly, with left over energy to enjoy leisuretime activities. It also helps to withstand stress and carry on, in circumstances where a physically unfit person could not continue.

Agricultural work is one of the highly physically demanding occupations. Farmers handle different heavy workloads, often in awkward postures and experiencing some work related musculo-skeletal disorder [1]. Some researchers has developed a perspective which identifies that farmers experience high rates of low back, shoulder, and upper extremity disorders [2]. According to Gangopadhyay *et al.* states that preadolescent agricultural workers suffered discomfort feeling (pain) in different parts of the body

especially low back, knee, shoulder, hand and neck region due to working in an awkward posture for prolonged periods of time in the agricultural field [3].

Farmers and agricultural workers are believed to be the healthier and have lower morbidity and mortality rates than non-farming rural and urban populations [4-8]. This fact has been reported possibly attributable to a healthier lifestyle, especially with respect to drinking and smoking habits, more intensive physical activity and a healthier diet followed by farmers compared to non-farming populations [4, 9, 10].

On the other hand, farming itself and farming related tasks entail significant hazards to the health and well-being of farmers. Although not well appreciated, farming is among the most hazardous of occupations [11]. Farmers work long hours in hazardous and physically demanding work environments [12]. Health impairments observed on farmers is a highly controversial issue and many studies have focused on agricultural work-related factors that may have a health impact [4, 11, 13-16]. Agricultural workers are exposed to a wide range of occupational hazards, such as ergonomic stress, sunlight, viruses, inorganic dust, pesticides and other chemicals [13]. All these exposures

have been investigated as possible risk factors for the reported adverse health effects in farmers including musculoskeletal disorders, respiratory diseases, injuries, cardiovascular diseases, pesticides poisoning and neurological dysfunction [12, 13, 17].

Moreover, stress in farm workers has been recently recognized as an important public health concern. Stressors inherent in farm work and lifestyle, such as uncertain and fluctuating economic prospects are associated with poor physical and mental health outcomes and result in deleterious effects on cognitive function, depression and high rates of suicides [13, 16, 18].

In India, about 20% of the labor force population is engaged in agriculture. While agriculture is one of the most promising sectors of the Indian economy, little interest has been demonstrated in studying health and safety of the farming population. Farmers in Greece are involved in all kind of agricultural tasks [19].

## MATERIALS AND METHODS

### Source of data

The present study was a cross-sectional study, conducted in the different villages of an agricultural area in Bankura district of West Bengal, where about 60% of the total population deals with various cultivations. It is worth noting that the majority of the farming population in this area is indigenous and non-immigrants. The study population was acquired from 36 rural communities of the prefecture.

### Method of collection of data

#### Sample size

A statistically adequate sample of 45 male farmers and 45 non-farmers in the age group of 25-40 years who were performed their tasks according to their occupations was selected from a process organization.

#### Type of work

To participate in the study, the farmers had to be at least thirty years old, so as to satisfy the criterion of long term farming (at least 10-15 years of farming), given that farmers start dealing with intensive agricultural work at the age of 17-18 years old. Non-farmers had to live in the same communities, not to be occupationally involved in farming and to match the recruited farmers *per* sex and age category.

#### Anthropological parameters

Measurements of body weight and height and Body Mass Index were determined by using height and weight (Quetelet Index method). BMI was calculated from self regulated weight (converted from pounds to kilograms divided by height (converted from inches to meters) squared by Quetelet Index [20-22] and their

level of obesity was determined as per WHO scores (Table-1).

#### Measurement of pulse rate

Pulse rate is measured in right radial artery, after all participants had rested for at least 10 mins before and after exercise. Pulse rate was measured by using three fingers and for a complete minute in supine, in sitting and standing positions. The results were recorded as pulse rate per minute and then analyzed.

#### Measurement of Blood pressure

Blood pressure was measured by the auscultatory method in the right arm in supine, sitting and standing position by using a mercury sphygmomanometer with a cuff of 12 cm. widths. All the subjects were made to rest for at least 10 mins before taking the readings. The manometer cuff was snugly tied around the arm with tubing on the medial and the lower side. Systolic blood pressure was recorded to the nearest 2 mm of Hg at appearance of first korotkoff sound, and diastolic blood pressure was recorded to the nearest 2 mm of Hg at the disappearance of korotkoff sound. Systolic and diastolic blood pressure was recorded first in the supine position and then standing position, with cuff tied to the arm. Reading was taken in all 3 positions and was analyzed [23].

#### Measurement of Physical Fitness Index (PFI)

PFI was measured by modified Harvard Step test method. PFI was calculated by measuring heart rate after performing the Harvard step test (HST) developed by Brouha *et al.* in the Harvard Fatigue Laboratories using long form PFI equation [24]. But, following modified HST under Indian condition, using stool of 51 cm high stepping up and down with a rate of 30 cycles/min for 3 minutes or up to exhaustion. Exhaustion is defined as when the subject cannot maintain the stepping rate for 15 seconds [25, 26]. The recovery pulse was counted at 1 to 1.5, 2 to 2.5 and 3 to 3.5 minutes of recovery.

#### Statistical analysis

The questionnaires were filled through interviews with both of the groups. For data analysis, SPSS 18, t-test and Chi-Square were used; the meaningful level was lower than 0.05 in almost all the analyses.

## RESULTS

In this study it was observed that differences in the educational status among farmers and non-farmers. Usually farmers showing extremely low percentages, especially regarding secondary or higher degrees. Furthermore, higher income levels were observed in farmers than non-farmers. The percentage of heavy smokers was higher in farmers than non-farmers.

Table 2 shows the mean age of farmer was  $33 \pm 7$  years and mean age of non-farmers were  $32 \pm 5$ . Mean height, body mass and BMI shows that both farmers and non-farmers BMI score were in healthy level as per WHO scores, and their prevalence of obesity is very minimal for both. The anthropometric measurements between both farmers and non-farmers did not show any significant change.

Mean and standard deviation of heart rate and blood pressure were recorded and it was observed that diastolic blood pressure for non-farmers were more than farmers (Table 3) (Figure-1).

Physical fitness index and their recovery period duration were measured among both farmers and non-farmers. It was found that farmers are physically more fit

than the non-farmers may be due to their activities which are related with farming (Table 4). The PFI score was also statistically significant ( $p < 0.001$ ).

Farmers reported that they have more musculoskeletal symptoms than non-farmers especially in the elbow, fingers, waist regions and also sometimes in finger joints (Table 5). The differences were significant for low back and hip symptoms. The farmers reported significantly more workload, more vibrations, more heavy lifting, more difficult working positions, longer work and sleep hours, less leisure-time physical activity than the non-farmers. They also had significantly higher total muscle strength and arm strength, and had a significantly higher physical work capacity.

**Table 1: WHO Classification of BMI (Sources from WHO)**

BMI	Category
<18.5	Underweight
18.5-24.9	Healthy
25-29.9	Overweight
30-39.9	Obese
>40	Morbid obese

**Table 2: Comparison of BMI among Farmers and non-farmers (Mean and standard deviation (S.D.) of anthropometric parameter)**

Parameters	Farmers	Non-Farmers
Sample Size	45	45
Age	$33 \pm 7$	$32 \pm 5$
Sex	Male	Male
Mean Height (cm)	$160.6 \pm 7.08$	$162.5 \pm 8.55$
Mean Weight (kg)	$58.37 \pm 8.89$	$60.6 \pm 10.15$
Mean BMI	$22.61 \pm 2.89$	$22.78 \pm 3.0$

\*Values were in (Mean  $\pm$  SEM)

**Table 3: Comparison of cardiovascular parameters among Farmers and non-farmers (Mean and standard deviation (S.D.) of anthropometric parameter)**

Parameters	Farmers (n=45) (Mean $\pm$ SEM)	Non-Farmers (n=45) (Mean $\pm$ SEM)	p Value
HR (Beats/ min)	$82.22 \pm 10.62$	$84.46 \pm 18.06$	$p < 0.05$
SBP (mm of Hg )	$121.06 \pm 6.86$	$124.04 \pm 14.92$	$p < 0.05$
DBP (mm of Hg )	$78.62 \pm 3.56$	$80.07 \pm 6.83$	$p < 0.07$
PP (mm of Hg )	$42.22 \pm 5.42$	$43.73 \pm 8.27$	$p < 0.05$

\*P value  $< 0.05$  is considered significant


**Table 4: Comparison of physical fitness index among Farmers and non-farmers:**

Parameters	Farmers (n=45) (Mean $\pm$ SEM)	Non-Farmers (n=45) (Mean $\pm$ SEM)	P Value
Basal HR (Beats/ min)	$82.22 \pm 10.62$	$84.46 \pm 18.06$	$P < 0.005$
PFI score	$75.1 \pm 3.10$	$60.9 \pm 3.90$	$P < 0.001$
Recovery period (min)	$6 \pm 2$	$8 \pm 4$	$P < 0.005$

\*P value  $< 0.05$  is considered significant

**Table 5: Musculoskeletal symptoms among Farmers & Non-farmers**

Parameters	Farmers (n=45)	Non-Farmers (n=45)
Elbow	Yes	No
Wrist	No	No
Fingers	Yes	No
Waist region	Yes	No
Knee-joint	No	Yes
Legs finger	No	No
Low back pain	Yes	No



**Fig-1: Graphical representation of mean cardiovascular parameters among farmers and non-farmers**

### DISCUSSION AND CONCLUSION

The present study gives evidence of several clinical and neurobehavioral alterations among farmers who have been involved in cultivation for a long time-period, indicating their susceptibility to certain impairments of their health status in comparison with non-farmers in the same area. Based on the participants' statements regarding their health problems, farmers mention more health issues than non-farmers at a statistically significant level ( $p < 0.005$ ). Hypertension and other cardiovascular disorders occur more frequently among farmers in comparison to non-farmers. It may be due to their smoking habits (such as bidi, and alcohol consumption like haria).

Farmers and non-farmers both have several musculoskeletal disorders and joint pains such as waist region, low back pain. It may be due to involvement of such body parts most actively during farming. But farmers have some more pains which the non-farmers don't have. Such type of pains is elbow pain and pain in fingers.

This is a pilot study, which makes an attempt to link the health & farm production in a framework of agricultural household model. The profit function approach has been adopted to analyze the issue econometrically. The analysis is, however, done both with the help of descriptive tables and econometric tools. Health status is found to be an important determinant of farm profit.

### REFERENCES

1. Kumudini, G., & Hasegawa, T. (2009). Workload and awkward posture problems among small-scale strawberry farmers in Japan. *Journal of human ergology*, 38(2), 81-8.
2. Davis, K. G., & Kotowski, S. E. (2007). Understanding the ergonomic risk for musculoskeletal disorders in the United States agricultural sector. *American Journal of industrial medicine*, 50(7):501-11.
3. Gangopadhyay, S., Das, B., Das, T., & Ghoshal, G. (2006). The prevalence of musculoskeletal disorders among pre-adolescent agricultural workers of West Bengal, India. *Ergonomics SA: Journal of the Ergonomics Society of South Africa*, 18(1), 14-21.
4. Stiernström, E. L., Holmberg, S., Thelin, A., & Svärdsudd, K. (1998). Reported health status among farmers and non-farmers in nine rural districts. *J. Occup. Environ. Med.*, 40, 917-924.
5. Fleming, L. E., Bean, J. A., Rudolph, M., & Hamilton, K. (1999). Mortality in a cohort of licensed pesticide applicators in Florida. *Occup. Environ. Med.*, 56, 14-21.
6. Stiernström, E. L., Holmberg, S., Thelin, A., & Svärdsudd, K. (2001). A prospective study of morbidity and mortality rates among farmers and rural and urban nonfarmers. *J. Clin. Epidemiol.*, 54, 121-126.
7. Wang, Y., Hwang, S. A., Lewis-Michl, E. L., Fitzgerald, E. F., & Stark, A. D. (2003). Mortality among a cohort of female farm residents in New York State. *Arch. Environ. Health*, 58, 642-648.
8. Thelin, N., Holmberg, S., Nettelbladt, P., & Thelin, A. (2009). Mortality and morbidity among farmers,

- nonfarming rural men, and urban referents: A prospective population-based study. *Int. J. Occup. Environ. Health*, 15, 21–28.
9. Pomrehn, P. R., Wallace, R. B., & Burmeister, L. F. (1982). Ischemic heart disease mortality in Iowa farmers: The influence of life-style. *JAMA*, 248, 1073–1076.
  10. Garcia-Palmieri, M. R., Sorlie, P. D., Havlik, R. J., Costas, R., & Cruz-Vidal, M. (1988). Urban-rural differences in 12 year coronary heart disease mortality: The Puerto-Rico heart health program. *J. Clin. Epidemiol.*, 41, 285–292.
  11. Frank, A. L., McKnight, R., Kirkhorn, S. R., & Gunderson, P. (2004). Issues of agricultural safety and health. *Annu. Rev. Publ. Health*, 25, 225–245.
  12. Brumby, S., Chandrasekara, A., McCoombe, S., Kremer, P., & Lewandowski, P. (2012). Cardiovascular risk factors and psychological distress in Australian farming communities. *Aust. J. Rural Health*, 20, 131–137.
  13. Lee, W. J., Cha, E. S., & Moon, E. K. (2010). Disease prevalence and mortality among agricultural workers in Korea. *J. Korean Med. Sci.*, 25, S112–S118.
  14. Waggoner, J. K., Kullman, G. J., Henneberger, P. K., Umbach, D. M., Blair, A., Alavanja, M. C., Kamel, F., Lynch, C. F., Knott, C., London, S. J., Hines, C. J., Thomas, K. W., Sandler, D. P., Lubin, J. H., Beane Freeman, L. E., & Hoppin, J. A. (2011). Mortality in the agricultural health study, 1993–2007. *Am. J. Epidemiol.*, 173, 71–83.
  15. Arcury, T., & Quandt, S. A. Living and working safely: Challenges for migrant and seasonal farmworkers. *N. C. Med. J.*, 72, 466–470.
  16. Nguyen, H. T., Quandt, S. A., Grzywacz, J. G., Chen, H., Galván, L., Kitner-Triolo, M. H., & Arcury, T.A. (2012). Stress and cognitive function in Latino farmworkers. *Am. J. Ind. Med.*, 55, 707–713.
  17. Rural, Regional and Remote Health Indicators of Health Status and Determinants of Health; Rural Health Series No. 9. Cat. No. PHE 97; Australian Institute of Health and Welfare (AIHW): Canberra, Australia, 2008. Available online: <http://www.aihw.gov.au/publication-detail/?id=6442468076> (accessed on 18 January 2013).
  18. Hounsome, B., Edwards, R.T., Hounsome, N., & Edwards-Jones, G. (2012). Psychological morbidity of farmers and non-farming population: Results from a UK survey. *Community Ment. Health J.*, 48, 503–510.
  19. Hellenic Republic. National Statistical Service of Greece. Available online: [http://www.statistics.gr/portal/page/portal/ESYE/PA GE-themes?p\\_param=A1604](http://www.statistics.gr/portal/page/portal/ESYE/PA GE-themes?p_param=A1604) (accessed on 18 January 2013).
  20. Garrow, J. S., & Webster, J. D. (1985). Quelet's index (W/H<sup>2</sup>) As a measure of fatness. *Int J of Obesity*, 9(2), 147-53.
  21. Panel, N. O. E. I. E. (1998). on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults—the evidence report. *Obes Res*, 6(Suppl 2), 51S-209S.
  22. World Health Organization. Obesity: Preventing and managing the Global Epidemic. Report of a WHO Consultation of Obesity. Geneva, 3-5 June 1997.
  23. He, J., Klag, M. J., Whelton, P. K., Chen, J. U., Qian, M. C., & He, G. Q. (1994). Body mass and blood pressure in a lean population in Southwest China. *Am J Epidemiol.*, 139, 380-9.
  24. Brouha, I., Health, C. W., & Gray, B. (1943). A step test simple method of measuring physical fitness for hard muscular work in adult men. *Rev Canadian Biol.*, 2, 86.
  25. Ryhming, I. (1953). A modified Harvard Step Test for Evaluation of Physical Fitness. *Arbeitsphysiologie*, 15(3), 235-50.
  26. Monotoye, H. J. (1953). The Harvard Step Test and Work Ca-pacity. *Rev Can Biol.*, 11(5), 491-9.