

Research Article

Cymbopogon citratus Stapf (DC) extracts alleviate gasoline vapour-induced metabolic disorders and cardiovascular disease risk in rats

Dr. Christopher E. Ekpenyong¹, Samson A. Oyebadejo²

¹Department of Physiology, Faculty of Basic Medical Sciences, University of Uyo, Nigeria

²Derindam Research Institute of Biotechnology, Uyo, Akwa Ibom State, Nigeria

*Corresponding Author:

Dr. Christopher E. Ekpenyong

Email: chrisvon200@yahoo.com

Abstract: Alternative therapies for the alleviation of chemical/xenobiotic-induced metabolic aberrations and risk factors for cardiovascular disease have been inadequately explored, despite their rich cardiovascular protective bioactive constituents. The aim of the present study was to accurately evaluate the effect of *Cymbopogon citratus* extract on gasoline-induced oxidative stress, hyperglycaemia, hyperuricemia, dyslipidaemia, and atherogenic index of plasma (AIP) in rats. Seventy-two female Wistar albino rats were segregated into six groups (n = 12 per group). The rats in group 1 served as controls, while animals in group 2 were exposed to gasoline vapour (GV) alone for 35 days. Rats in groups 3, 4, and 5 were exposed to GV and co-administered 500 mg/kg, 1000 mg/kg, and 1500 mg/kg *C. citratus* leaf extract, respectively, once daily for 14 days. Rats in group 6 were exposed to GV and co-administered 200 mg/kg vitamin C for the same time period. The results showed that group 2 had significantly higher glycaemic levels, lipid parameters (TG, T-Chol, and LDL-C), uric acid (UA), MDA, and AIP and significantly lower ($p < 0.05$) HDL-C than group 1. Co-administration of *C. citratus* leaf extracts in groups 3, 4, and 5 and vitamin C to group 6 significantly lowered blood glucose levels, UA, lipid parameters MDA and AIP and simultaneously increased serum HDL-C in a dose-dependent manner. *C. citratus* leaf extract has an ameliorative effect against GV-induced oxidative stress, hyperglycaemia, hyperuricaemia, dyslipidaemia, and AIP, probably mediated through its varied bio – constituents and their activities.

Keywords: lemongrass extract, lipid peroxidation, atherosclerosis, cardiovascular risk, rat

INTRODUCTION

Gasoline is a colourless to pale brown or pink liquid. It is produced by distillation, cracking, and reforming of crude oil. Gasoline has wide domestic and industrial applications, including as a fuel in spark ignition and internal combustion engines in automobiles, trucks, and aircraft. It is a volatile liquid that evaporates easily into the atmosphere, constituting a significant environmental pollutant and hazard to the general population and in particular, to those who are occupationally exposed [1].

Acute and chronic exposure to gasoline compounds is associated with several systemic adverse health effects, including abnormalities in serum blood sugar levels, and lipid parameters (dyslipidaemia), leading to an increased risk of atherosclerotic cardiovascular diseases [2].

Available data indicate that dyslipidaemia constitutes a considerable risk factor for coronary artery disease (CAD). Dyslipidaemia is characterized by significant increases in serum levels of low-density lipoprotein cholesterol (LDL-C), triglyceride (TG), and total cholesterol (T-Chol) and low levels of high-

density cholesterol (HDL-C). Previous studies have shown that exposure to gasoline may be associated with dyslipidaemia, and thus may constitute a significant risk factor for atherosclerosis and cardiovascular diseases [2]. Several studies [3] have indicated that exposure to gasoline vapour (GV) leads to significant increases in serum levels of TG, T-Chol, LDL-C, and VLDC, and a significant decrease in serum HDL-C, as well as a corresponding increase in the TG/HDL-C ratio, otherwise known as the atherogenic index of plasma (AIP). AIP is strongly associated with increased atherosclerotic cardiovascular disease risk in the general population. In fact, it is the most useful index for predicting and quantifying coronary artery disease (CAD) risk and morbidity and mortality in those affected [4]. The pathophysiological mechanisms underlying gasoline-induced dyslipidaemia have been postulated to include induction of oxidative stress (OS) [5], and increased serum levels of C-reactive protein [6]. OS is associated with the generation of free radicals, and induction of oxidative changes in low density lipoprotein cholesterol (LDL-C) particles that initiate and promote atherosclerotic changes. Current understanding supports the use of drugs with anti-

oxidative as well as anti lipid effects to mitigate gasoline-induced dyslipidaemia and consequences.

The use of synthetic lipid lowering drugs, such as fibrates and statins, has been very effective; however, they are associated with adverse health effects, such as myopathy [7], hepatotoxicity [8], and drug-drug interactions [9], which limit their use. Furthermore, long-term use of fibrates in dyslipidaemic patients is associated with increased plasma selenium [10]. High selenium levels (above the physiological limit) have been linked to hypertension [11], insulin resistance, obesity [12, 13, 14], hyperglycaemia [15], and diabetes mellitus. In addition, in some poor communities and developing countries, these drugs may not be accessible and/or affordable. Furthermore, synthetic anti-lipid and antioxidant drugs may not address the pleotropic effects of gasoline metabolite-induced dyslipidaemia, and complications such as hyperglycaemia and atherosclerotic cardiovascular disease, mandating the use of multiple drug therapies, which are associated with additional complications. Often, patients who are dissatisfied with their current treatments will seek alternative treatments [16]. Therefore, despite the large number of modern synthetic drugs, people see using complementary and alternative medicine (CAM) to prevent and treat illnesses as an attractive proposition. Thus, the search for natural products with bioactive constituents that can address health concerns is currently garnering increased attention. Many patients, healthcare providers, and environmental and biochemical toxicologists are fervently searching for CAM options to treat metabolic maladies. In one such search, Sirotkin et al. [17] found that medicinal plants can regulate ovarian cell function and alleviate the toxic effects of gasoline (benzene, toluene, ethylbenzene, and xylene [BTEX]) compounds on ovarian cells. Several reports [18, 19, 20] have demonstrated the feasibility of using plant products to treat hyperglycaemia, dyslipidaemia, and causative factors. Natural plant products are known to contain several bioactive constituents that can exert multiple anti-dyslipidaemic/anti-atherosclerotic and hypoglycaemic actions, such as antioxidative stress [21], radical scavenging [22], anti-inflammatory, and immunomodulating [23] activities, as well modify the genes of target cells and hormone secretion [24]. One such plant that has been explored for its anti-dyslipidaemic, hypoglycaemic, and antioxidant effects is *Cymbopogon citratus* (*C. citratus*) [20].

C. citratus (lemongrass) is a perennial tropical plant belonging to the Poaceae family that is known to have numerous bioactive constituents with the potential to prevent and treat various diseases through their antioxidant, anti-inflammatory, anti-protozoal, anti-fungal, anti-carcinogenic, anti-platelet aggregation, anti-bacterial, anti-tussive, anti-diabetic, and anti-malarial effects. It has also been used to treat

gastrointestinal disturbances, anxiety, flu, fever, pneumonia, and hypertension. However, whether *C. citratus* extract can alleviate gasoline-induced CVD risk factors remains undocumented. In view of the National and International recommendation to increase fruit and vegetable intake, there is the need to evaluate the relation between intake of these foods and CVD protective effect [25]. Therefore, the aim of the present study was to evaluate the effect of *C. citratus* leaf extract on GV-induced hyperglycaemia, hyperuricaemia, dyslipidaemia and AIP in rats.

MATERIALS AND METHODS

Selection and Segregation of Animals

The experimental animals were 72 matured female Wister albino rats weighing 180-200g. The animals were obtained from the animal house at Faculty of Basic Medical sciences, University of Uyo, Nigeria. They were acclimatized for one week before starting the experiment. Each animal was housed in a standard wooden cage with wood shavings as bedding, which was regularly replaced. The experiment was performed under standard laboratory conditions (at room temperature [$28 \pm 8^\circ\text{C}$], 45% humidity, with a 12-h light/dark cycle). They were randomly divided into six groups with 12 rats per group (n=12 per group): rats in group 1 (G1) served as the control (unexposed), rats in group2 (G2) served as exposed control, while rats in groups 3,4,5 and 6 served as test groups.

All research protocols were performed at the University of Uyo according to Nigerian and international laws (Revised Helsinki Declaration, 2008) governing the acceptable use of laboratory animals.

Collection and Preparation of *C. citratus* Leaf Extracts

A day prior to utilization, fresh *C. citratus* leaves were obtained from an agricultural farm in Uyo, Akwa Ibom State Nigeria. Identification and authentication was performed by a taxonomist (ID UHH 4686/Uyo) in the Department of Botany at the University of Uyo. The leaves were rinsed, sundried, and pulverized into powder using an electric blender to provide 400 g of material. The powder was soaked with 4 L of hot water in a conical flask and allowed to stand for approximately 10 h. After filtering the solution through Whatman No. 2 filter paper, the filtrate was evaporated to dryness by heating in a water bath at 40°C . The final solid extract was weighed with an electric balance (ACS-2E14; Surgifriend Medicals, Ltd., England), with a total yield of 30%. The prepared extract was stored in glass bottles at 4°C and was dissolved in physiological saline at 100 mg/mL. The phytochemical analysis of the leaf extracts was carried out using standard procedures to determine the levels of tannins, phenolics, saponins, alkaloids, deoxysugars, and anthraquinones [26, 27].

The median lethal dose (LD₅₀) was determined using standard methods [28]. From the LD₅₀ (5000 mg/kg), the low, medium, and high doses were calculated using a standard formula as 500 mg/kg, 1000 mg/kg, and 1500 mg/kg, respectively.

Exposure to GV

The animals in the test groups were exposed to unleaded gasoline purchased from a Nigerian National Petroleum Cooperation (NNPC) refuelling station on Itam-Ikot Ekpene Road in Uyo, Nigeria. The rats in the exposed groups were housed in their cages and exposed to GV in an exposure chamber (80 × 60 × 100 cm). Rats in the unexposed control group were kept in a GV-free section of the experimental house. Two calibrated 500-mL beakers each containing 100 mL of petrol were put in the modified chamber where the exposed groups were placed, and then the rats were allowed to inhale the GV in the chamber for 6 h (9 am–3 pm) daily for 35 consecutive days. At the end of the exposure period, the gasoline was removed; the initial and final volumes were recorded before and after exposure, respectively. The daily difference in volume was used to estimate the relative vapour exposure. The average exposure was approximately 80 mL/day.

Experimental treatment

Rats in groups 3, 4, 5 and 6 were in addition to being exposed to GV, co-treated with 500mg/kg, 1000mg/kg and 1500mg/kg of *C. Citrates* extracts and 200mg/kg of vitamin C respectively. All animals were fed normal rat pellets and allowed free access to food and water throughout the experimental period. The *C. citratus* leaf extracts and water-solubilized vitamin C were administered by oral gavage (using an intragastric syringe) for the final 14 days of the 35-day GV exposure.

Serum bio-chemical Analysis

Serum glucose level and lipid sub-fractions including total cholesterol (T.chol), Low density lipoprotein cholesterol (LDL-C), high density lipoprotein cholesterol (HDL-C) and triglyceride (TG) were determined using a multi-channel automated system (Lipid pro TM, Model KM-001A: Info Pia Co, Ltd. South Korea). Serum uric acid (UA) levels were measured using a multi-channel automated analyser (SYNCHRON, LOS ANGELES, CA).

Statistical Analysis

Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS), version 20.0.

Data obtained were analysed using descriptive statistics and reported as the mean ± standard deviation (SD). Analysis of variance (ANOVA) was also used. Duncan's multiple range test was used to compare results between groups, determine the direction of significance, and to analyse the effect of the *C. citratus* leaf extracts on the reproductive toxicity of GV. Differences with *p* values less than 0.05 were considered statistically significant.

RESULTS

Phytochemical analysis of *C. citratus* leaf extract showed the presence of moderate concentrations of tannins, saponins, flavonoids, and phenols and low levels of anthraquinone. Table 1 shows the ameliorative effect of *C. citratus* extract on serum lipid profile markers of GV-induced dyslipidaemic models in rats.

Figures 1, 2, 3 and 4 show the ameliorative effects of *C. citratus* leaf extracts on serum glucose, malondialdehyde, uric acid and atherogenic index of plasma (AIP) in GV-induced hyperglycaemic, lipid peroxidation, hyperuricaemic and dyslipidaemic models in rats respectively.

Table 1: The ameliorative effect of *C. citratus* extracts on serum lipid profile markers of gasoline vapour-induced dyslipidemic models in rats.

Groups	T-Chol (mg/dl)	TG (mg/dl)	HDL (mg/dl)	LDL (mg/dl)	TL (mg/dl)
1	67.50 ± 4.84 ^b	43.33 ± 7.03 ^a	29.00 ± 3.27 ^b	43.17 ± 14.18 ^a	91.83 ± 8.89 ^b
2	72.50 ± 3.54 ^c	69.00 ± 4.24 ^b	18.00 ± 3.22 ^a	63.50 ± 10.61 ^a	105.50 ± 3.54 ^c
3	72.33 ± 3.56 ^c	52.33 ± 7.31 ^c	20.10 ± 1.27 ^a	48.33 ± 9.85 ^a	97.33 ± 2.88 ^b
4	65.33 ± 2.07 ^b	56.50 ± 3.44 ^c	23.25 ± 3.44 ^a	43.33 ± 7.94 ^a	85.83 ± 4.88 ^a
5	55.67 ± 4.32 ^a	56.17 ± 4.79 ^c	25.32 ± 1.59 ^b	33.33 ± 15.19 ^a	92.33 ± 7.53 ^b
6	57.50 ± 3.62 ^a	47.50 ± 3.02 ^b	27.23 ± 2.88 ^b	34.50 ± 5.01 ^a	77.33 ± 12.24 ^a

The different superscripts are significantly different from control at 5% (*p* < 0.05)




Fig-1: Effect of *C. citractus* Extract (CCE) and Vitamin C Supplementation on GV-induced Hyperglycaemia (mg/dl).




Fig-2: Effect of *C. citractus* Extract (CCE) and Vitamin C Supplementation on GV-induced increase Serum Malondialdehyde (mmol/L).




Fig-3: Effect of *C. citractus* Extract (CCE) and Vitamin C Supplementation on GV-induced increase Serum Uric Acid Concentration (mg/dl).




Fig- 4: Effect of *C. citractus* Extract (CCE) and Vitamin C Supplementation on GV-induced increase Atherogenic Index of Plasma (AIP).

DISCUSSION

The results from the present study showed that exposure to GV alone significantly increased serum glucose, UA, TG, LDL-C, T-chol, MDA, and AIP, and significantly decreased serum HDL-C.

When rats in group 3, group 4, and group 5 were co-administered 500 mg/kg, 1000 mg/kg, and 1500 mg/kg of *C. citratus* leaf extract, respectively, we observed significant dose-dependent decreases in the aforementioned lipid parameters, blood glucose, UA, MDA, and AIP, and a significant increase in HDL-C. These changes were similar to those observed in rats

(group 6) co-administered vitamin C (an antioxidant reference). The present results highlight the etiopathogenic role of oxidative stress in GV-induced dyslipidaemia, hyperglycaemia, hyperuricaemia and atherosclerosis, as was previously documented [29, 3, 2].

Numerous established studies have confirmed that GV is readily absorbed when inhaled, and after a series of biotransformative and toxicokinetic processes, the GV is transformed into reactive metabolites, leading to the generation of toxic reactive oxygen species (ROS), such as superoxide (O⁻), hydroxyl radical (OH⁻),

hydrogen peroxide (H₂O₂), peroxy hydroxyl, and organic hydrogen radical [30, 2]. These reactive oxygen intermediates are known to induce oxidative stress that is a critical pathophysiological step in GV-induced abnormalities in serum lipid parameters (dyslipidaemia) [31], such as significantly increased levels of serum glucose, LDL-C, VLDL-C, TG, and T-chol and a simultaneous decrease in HDL-C, as was observed in the present study. The mechanisms underlying GV-induced dyslipidaemia and subsequent atherosclerosis may include, in part, the inactivation of metabolic enzymes and damage to important cellular components and oxidation of nucleic acids [30]. These are associated with disordered protein, nucleic acid, carbohydrate, and lipid metabolism, leading to alterations in intracellular and extracellular homeostasis and subsequent loss of cell integrity, enzyme function, and genomic stability [32]. For example, enhanced lipid peroxidation [31], including the oxidation of LDL-C, is a necessary step in the development of atherogenesis [33]. GV-induced dyslipidaemia has also been thought to be due to GV-induced liver toxicity, since the liver functions in lipid metabolism and homeostasis [2]. Interestingly, these processes can be mitigated by the use of nutritional antioxidants [30], particularly green leafy vegetables and vitamin-rich fruits [34, 35] such as *Cymbopogon citratus* [36, 37, 38, 20], which is known to decrease the toxic events of lipid peroxidation in both serum and heart tissue [39]. The significant increase in serum levels of MDA in rats exposed to GV alone (G2) indicates increased lipid peroxidation, a reaction between free radicals and polyunsaturated fatty acids in the cell membrane [31] following oxidative stress.

The significant dose-dependent decreases in some lipid sub-fractions (LDL-C, TG, and T-chol), the lipid peroxidation marker MDA, and CAD-risk indicator AIP, and the simultaneous increase in the good lipid sub-fraction (HDL-C) in groups co-administered *C. citratus* leaf extracts or vitamin C compared with the corresponding levels in the control group confirmed the antioxidant effect of *C. citratus* leaf extract, which is likely due to its rich and varied antioxidant constituents and their activities, as was reported in previous studies [40]. Several of the *C. citratus* constituents, including some phytochemicals (flavonoids, tannins, polyphenols, alkaloid, and saponins), vitamins (A, C, and E), and minerals and trace elements (zinc, selenium, magnesium, and copper) [20], have documented antioxidant activities that likely contribute to their anti-dyslipidaemic, hypouricaemic, and hypoglycaemic effects [41,42,43]. In our earlier study [20] on the effect of *C. citratus* leaf extract on AIP in diabetic dyslipidaemic animals, administration of *C. citratus* extract dose-dependently abated the rise in serum levels of T-chol, LDL-C, VLDL-C, β -hydroxyl β -methyl glutaryl-CoA (HMGCoA) reductase, and AIP and simultaneously increased HDL-C. These results support those of a previous study by Adeneye


and Agbaje [44], which was designed to investigate the hypoglycaemic and hypolipidaemic effects of *C. citratus*; the authors found that *C. citratus* leaf extract lowered fasting plasma glucose and lipid parameters dose-dependently, while raising plasma HDL-C. Accordingly, Ewenighi et al. [45] found that a four-week treatment with *C. citratus* leaf extracts mitigated the increases levels in serum TG, T-chol, and LDL-C in diabetic animals compared with the levels in the untreated group. In a parallel study, Agbafor et al. [46] confirmed the anti-dyslipidaemic effect of *C. citratus* leaf extract, which was traceable to its phytochemical constituents. In addition, the presence/effect of these bioactive antioxidants in *C. citratus* leaf have been used to abate drug/xenobiotic-induced oxidative stress and adverse health endpoints, leading to increased levels of the key antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx) and simultaneously decreasing the level of the lipid peroxidation marker MDA [40]. Koh et al. [47] found that prophylactic treatment with *C. citratus* dose-dependently ameliorated carbon tetrachloride-mediated hepatic oxidative damage in animals. This effect was attributable to the antioxidant and free radical-scavenging properties of the extract. In a parallel study, Ojo et al. [36] reported the antioxidative properties of *C. citratus* in paracetamol-induced oxidative stress in animals. Likewise, a study by Olorunsanya et al. [37] reported that *C. citratus* inhibited lipid oxidation in raw pork patties under refrigeration. Besides the antioxidant effect of some *C. citratus* constituents, the essential oil (EO) constituents myrcene, geraniol, linalool limonene, β -ionone, and citral (α and β) have been shown to possess antioxidant activities in human cells [48] and cholesterol-reducing activities in mice [49,50]. Soares et al. [51] compared the antioxidant capacity of *C. citratus* EO with that of a synthetic antioxidant butylated hydroxytoluene (BHT), and found that *C. citratus* EO has greater antioxidant activity than the methanolic extracts and similar antioxidant activity to that of BHT. Several other authors have reported the antioxidant activity of *C. citratus* EO [52,53,54,55] and evaluated the application of *C. citratus* EO as a replacement for the commercial antioxidant BHT in food preservation. In addition, citral, an EO constituent, has been shown to lower serum cholesterol and inhibit lipogenesis by post-transcriptionally inhibiting HMG-CoA reductase, an enzyme that catalyzes the rate-limiting step of cholesterol biosynthesis and a major cellular target of many anti-dyslipidaemic drugs [56]. These rich antioxidant constituents of *C. citratus* leaf extract not only prevent direct damage due to gasoline compounds but also may inhibit the signalling pathways leading to inflammation as a result of its anti-inflammatory actions [57,58], thereby inhibiting the inflammatory component of gasoline-induced atherosclerosis. Support for this view comes from the observation by Gbenou et al [58], that *C. citratus* EO displayed significant dose dependent anti-inflammatory,

analgesic and antipyretic effects when orally administered to rats. Furthermore, numerous established studies have confirmed the hypoglycaemic effect of *C. citratus* phytoactive constituents [59,20].

In all likelihood, the hypoglycaemic effects of these constituents are due, at least in part, to their antioxidative activities. For instance, tannic acid stimulates glucose transport and induces GLUT translocation [60]. Polyphenols stimulate insulin release from the beta cells of the pancreas and inhibit α -glucosidase [61]. In related study, Bharti et al [62] showed that *C. citratus* EO significantly decreased glycosylated haemoglobin (HbA1c) and insulin levels as well as insulin resistance and oxidative stress markers in diabetic rats, while simultaneously increasing β -cell mass and islet number.

Furthermore, exposure of the experimental animals to GV alone caused significant increase in serum level of uric acid (UA), suggesting the hyper-uricaemic effect of GV. Hyper-uricemia is an independent risk factor for essential hypertension, metabolic syndrome and cardiovascular diseases [63,64]. Other diseases associated with hyper-uricemia which could have directly or indirectly impacted cardiovascular endpoints include autoimmune, inflammatory, endocrine and metastatic disorders [4]. Interestingly, when the GV exposed animals were co-treated with *C. citratus* extracts, a dose-dependent significant ($p < 0.05$) decreases in serum levels of UA were observed, and were similar to the effect of vitamin C, a reference antioxidant and hypouricaemic agent [4].

These findings are similar to those previously observed by Mirghani et al [65] and confirm the hypouricaemic effect of *C. citratus* leaf extract. In their overall assessment of the xanthine oxidase (XO) inhibitory potential of *C. citratus* extract, Mirghani et al [65] concluded that *C. citratus* has a high potentiality to be an alternative source of antigout drug, as it has a potential compound containing XO inhibitor. This idea is supported by the detection of high levels of polyphenol-compounds (tannins, flavonoids and phenols), electrolytes (potassium (K^+), Magnesium (Mg^{2+})), vitamin C in *C. citratus* leaf extracts in the present and previous studies [4]. The hypouricaemic effects of these compounds have been extensively documented [4]. For instance, polyphenol compounds have been shown to possess high XO inhibitory activity, and their ability to decrease serum UA levels has also been reported [66]. XO is an enzyme that catalyzes the hydroxylation of hypoxanthine to xanthine and UA in purine metabolic pathway [67] according to the following reaction:



Polyphenol compounds such as flavonoids block these reactions by competitively binding to the specific active site of XO altering its stereochemistry and hence biochemical effects [4]. This inhibitory action is enhanced by the presence of Olefins at carbon 2 (C2) and Carbon 3 (C3) that maintains the planar structure of flavonoids, the hydroxyl moiety at C7 and C5, and the carbonyl group at C4 in the flavonoid structure.

The presence of these functional groups has been shown to enhance favourable hydrogen bonds and electrostatic interactions between flavonoids and the XO active site, which are the prerequisites for the potent inhibitory action of flavonoids on XO [68]. Additionally, the increase in serum UA levels in the GV alone group could, in part, be attributed to the effect of GV on renal endpoints, impairing excretion as documented previously [69,4]. The kidneys excrete about 70% of UA generated daily to maintain homeostasis [70], therefore impaired renal function can lead to increase serum UA level as observed previously following exposure to GV [69,71]. The observed improvement in the groups (G3, G4, G5) and G6 co-treated with various doses of *C. Citrates* extracts and vitamin C respectively, could partly be due to the ameliorative effect of *C. citratus* leaf extracts and vitamin C on GV induced renal function impairment as reported in previous studies [72,73].

CONCLUSION

In conclusion, *C. citratus* leaf extract has the potential to abate GV-induced hyperglycaemia, hyperuricaemia, dyslipidaemia, and atherosclerotic cardiovascular disease risk. Its action is comparable to known synthetic antioxidants, and it may possess a wider spectrum of actions due to its numerous components with various activities that can address the pleotropic effects of GV-induced detrimental health effects.

Conflicts of Interests

The authors declare that they have no competing interest.

REFERENCES

1. El-Ghazaly, N., Ali, A., Dekinesh, S., Sedky, A., Kabeil, S. (2016) Hepatotoxicity of gasoline as an environmental pollutant on Albino Mice. *Athens J. Sci.* 30-40.
2. Sedky, A., El Sawy, H. (2015) Protective effect of vitamin C and E against gasoline vapours induced hematological and biochemical changes in male rats. *J. Sci. Res.* 7(3), 139-149.
3. Uboh, F.E., Akpanbiatu, M.I., Alozie, Y. 2008. Exposure to gasoline vapours: A potential risk factor for Atherosclerosis in male and female rats. *J. pharmacol. Toxicol.* 3(8), 600-609.

4. Ekpenyong, C.E., Daniel, N. 2015. Role of diets and dietary factors in the pathogenesis, management and prevention of abnormal uric acid levels. *Pharma. Nutr.* 3, 39-45.
5. Shehata, A.M., Yousef, O.M. 2010. Physiological studies on the risk factors responsible for atherosclerosis in rats. *Nature Sci.* 8(5), 144-151.
6. Ma, C.M., Lin, L.Y., Chen, H.W., Huang, L.C., Lli, J.F., Chuang, K.J. 2010. Volatile organic compounds exposure and cardiovascular effects in hair salons. *Occu. Med.* 60(8), 625-630.
7. Siqueire Carvalho, A.A., Poti Lima, O.W., Valiente, R.A. 2004. Statin and fibra associated myopathy. *Arq Neuro-Psiquiatr.* 62 (2-A), 252-261.
8. Parra, J.L., Reddy, K.R. 2003. Hepatotoxicity of hypolipidemic drugs. *Clin. liver dis.* 7, 415-433.
9. Trifiro, G., Corrao, S., Alacqua, M., Moretti, S., Tan, M., UVEC group et al. 2006. Interaction risk with proton pump inhibitors in general practice significant disagreement between different drugs related information sources. *Br. J. Clin. Pharmacol.* 62(5), 582-590.
10. Arnaud, J., Akbaraly, T.N., Hininger-favier, L., Berr, C., Roussel, A.M. 2009. Fibrates but not statins increase plasma selenium in dyslipidemic aged patient-The EVA study. *J. Trace Elem. Med Biol.* 23, 21-28.
11. Laclaustra, M., Navas-Acien, A., Stranges, S., Ordovas, J.M., Guallar, E. (2009) Serum Selenium Concentration and diabetes in U.S Adults National Health and Nutrition Examination Survey (NHANES) 2003-2004. *Environ. Health Perspectives*, 117(9), 1409-1413.
12. McClung, J.P., Roneker, C.A., Mu, U. et al. (2004) Development of insulin-resistance and obesity in mice over expressing cellular glutathione peroxidase. *Proc. Nat. L. Acad. Sci. USA.* 101, 8852-8857.
13. Li, X., Chen, H., Epstein, P.N. (2006) Metallothionein and catalase sensitize to diabetes in non-obese diabetic mice: reactive oxygen species may have a protective role in pancreatic beta-cells. *Diabetes*, 55, 1592-1604.
14. Wang, X.D., Vatamaniuk, M.Z., Wang, S.K., Roneker, C.A., Simmons, R.A., Lei, X.G. (2008) Molecular mechanisms for hyperinsulinemia induced by overproduction of selenium-dependent glutathione peroxidase in mice. *Diabetologia*, 5, 1515-1524.
15. Satyanarayana, S., Sekhar, J.R., Kumar, K.E. et al. (2006) Influence of Selenium (antioxidant) on gliclazide induced hypoglycaemia/and hyperglycaemia in normal/alloxan-induced diabetic rats. *Mol Cell Biochem*, 283, 123-127.
16. Soeken, K.L., Miller, S.A., Ernst, E. (2003) Herbal medicine for the treatment of rheumatoid arthritis: a systematic review. *Rheumatology*, 42, 652-659.
17. Sirotkin, A.V., Kadasi, A., Balazi, A., Bakova, Z., Harrath, A.H., Makarevich, A.V. et al. (2012) Influence of petrochemical industry environmental contaminants on animal ovarian cells. *J Microbiol Biotech Food Sci*, 2, 517-525.
18. Reach, K.I., Ernst, F. (1995) Garlic (*Allium sativum*) a potent medicinal plant. *Fortschr Med*, 113, 311-15.
19. Kaur, G., Meena, C. (2013) Evaluation of antihyperlipidemic potential of combinatorial extract of curcumin piperine and quercetin triton induced hyperlipidemia in rats. *Scin Int*, 1(3), 57.
20. Ekpenyong, C.E., Davies, K., Antai, E.E. (2014) Cymbopogon citratus stapf (DC) extract ameliorates atherogenic cardiovascular risk in diabetes-induced dyslipidemia in rats. *British J Med Medical Res*, 4(28), 4695-4709.
21. Keen, C.L., Holt, R.R., Oteiza, P.I., Fraga, C.G., Schmitz, H.H. (2005) Cocoa antioxidants and cardiovascular health. *Am J Clin Nutr*, 81, 2985-303.
22. Murota, K., Tarao, J. (2003) Antioxidative flavonoid quercetin: Implication of its intestinal absorption and metabolism. *Arch Biochem Biophys*, 417, 12-17.
23. Yamasaki, H. (1976) Pharmacology of sinomenine, an anti-rheumatic alkaloid from sinomenium acutum. *Acta Med Okayama*, 30, 1-20.
24. Satoh, H. (2014) Pharmacological effectiveness of the active phytochemicals contained in foods and herbs. *J Int Ethnopharmacol*, 3(4), 196-200.
25. WHO (1990) Report of a WHO Study Group. Diet, nutrition and the prevention of chronic disease World Health organ. *Tech Rep*, 191.
26. Trease, G.E., Evans, W.C. (1989) A Textbook of Pharmacognosy 13th ed., Balair, Tindall, London. pp. 100-101.
27. Sofowora, A. (1996) Medicinal plant and traditional medicine in Africa. Second ed., Spectrum Books, Ibadan, Nigeria, pp. 112.
28. Lorke, D. (1983) A new approach to practical acute toxicity testing. *Arch Toxicol*, 54, 275-287.
29. Uboh, F.E., Akpanbiatu, M.I., Ekaidem, I.S., Ebong, P.E., Umoh, I.B. (2007) Effect of inhalation exposure to gasoline on sex hormones profile in wistar albino rats. *Acta Endocrin 1-Bucharest*, 3, 23-30.
30. Rahman, T., Hosen, I., Towhidul Islam, M.M., Shekhar, H.U. (2012) Oxidative stress and human health Advances in Bioscience and Biotechnology. 3, 997-1019.
31. Odewabi, A.O., Ogundahunsi, O.A., Oyalowo, M., (2014) Effect of exposure to petroleum fumes on plasma antioxidant defense system in petrol attendants. *Br J Pharmacol Toxicol*, 5, 83-87.
32. Gille, J.J., Van Berkel, C.G., Joenje, H. (1994) Mutagenicity of metabolic oxygen radicals in mammalian cell cultures. *Carcinogenesis*, 15, 2695-2699.

33. Steinberg, D. (1997) Low density lipoprotein oxidation and its pathobiological significance. *J Biol Chem*, 272, 20963-20966.
34. Joshipura, K.J., Hu, B.F., Manson, J.E., Stampfer, M.J., Rimm, E.B., Speizer, F.E., Colditz, G., Ascherio, A., Rosner, B., Spiegelman, D., Willett, W.C. (2001) The Effect of Fruit and vegetable intake on Risk for Coronary Heart Disease. *Ann Intern Med*, 134(12), 1106-14.
35. Bazzano, L.A., He, J., Ogden, L.G., Loria, C.M., Vupputuri, S., Myers, L., Whelton, P.K. (2002) Fruit and vegetable intake and risk of cardiovascular disease in US adults; the first National Health and Nutritional Examination Survey. Epidemiologic Follow-up Study. *Am J Clin Nutr*, 76, 93-99.
36. Ojo, O. O., Kabutu, F. R., Bello, M., Babayo, U. (2006) Inhibition of Paracetamol-Induced Oxidative Stress in Rats by Extract of Lemongrass (*Cymbopogon citratus*) and Green Tea (*Camellia Sinensis*) in Rats. *African J Biotech*, 5, 1227 – 1232.
37. Olorunsanya, A.O., Olorunsanya, E.O., Bolu, S.A.O., Adejunmobi, C.T., Kayode, R.M. (2010) Effect of graded levels of lemongrass (*cymbopogon citratus*) on oxidative stability of raw or cooked pork patties. *Pak J Nutr*, 9(5), 467-470.
38. Pei, H.K., Mokhtar, R.A., Mohammed, I. (2012) Antioxidant potential of *cymbopogon citratus* extract: alleviation of carbon tetrachloride-induced hepatic oxidative stress and toxicity. *Hum Exp Toxicol*, 31(1), 81-91.
39. Gayathri, K., Jayachandran, K.S., Vasanthi, H.R., Rajamanickam, G.V. (2011) Cardio-protective effects of lemongrass as evidence by biochemical and histo-pathological changes in experimentally induced cardio-toxicity. *Human Exp Toxicol*, 30(8), 1073-1082.
40. Arhoghro, E.M., Kpomoh, D.E., Uwakwe, A.A. (2012) Curative potential of aqueous extract of lemon grass (*Cymbopogon Citratus*) on cisplatin-induced Hepatotoxicity in Albino Wister Rats. *J Phys Pharm Adv*, 2(8), 282-294.
41. Fuentes, J. L., Alonso, A., Cuetara, E., Vernhe, M., Alvarez, N., Sanchez-lamar, A., Liangstera, M. (2006) Usefulness of the SOS Chromotest in the study of medicinal plants as radioprotectors. *Int J Radiation Bio*, 82(5), 323-329.
42. Anderson, R. A., Polansky, M. M. (2002) Tea Enhances Insulin Activity. *J Agric Food Chem*, 50, 7182 – 6
43. Serrano, J., Puupponen-Pimia, R., Dauer, A., Aura, A., Saura-Calixto, F. (2009) Tannins: Current Knowledge of Food Sources, Intake, Bioavailability and Biological Effects. *Mol Nutr Food Res*, 53, S310-29
44. Adeneye, A. A., Abaje, E. O. (2007) Hypoglycemic and Hypolipidemic Effects of Fresh Leaf Aqueous Extract of *Cymbopogon Citratus* Stapf in Rats. *Journal of Ethnopharmacology*, 12 (3), 440-444.
45. Ewenighi, C.O., Dimkpa, U., Adejumo, B.I., Onyeanus, J.C., Onah, L.U., Ezeugwu, U. et al. (2013) Estimation of lipid profile and glucose level in alloxan-induced diabetic rats treated with *cymbopogon citratus* (Lemongrass). *J Exp Integr Med*, 3(3), 249-253.
46. Agbafor, K.N., Akubugwo, E.I. (2007) Hypocholesterolemic effects of ethanolic extract of fresh leaves of *cymbopogon citratus* (Lemongrass). *Afri J Biotechnol*, 6: 596-598.
47. Koh, P.H., Mokhtar, R.A., Iqbal, M. (2012) Antioxidant potential of *Cymbopogon citratus* extract: alleviation of carbon tetrachloride-induced hepatic oxidative stress and toxicity. *Hum Exp Toxicol*, 31, 81-91.
48. Saeio, K., Caiyana, W., Okonogi, S. (2011) Antityrosinase and antioxidant activities of essential oils of edible Thai plants. *Drug Discoveries Therapeutics*, 5, 144-149.
49. Costa, C.A., Bidinotto, L.T., Takahira, R.K., Rodrigues, M.A., Barbisan, L.F. (2011) Cholesterol reduction and lack of genotoxic or toxic effects in mice after repeated 21 days oral intake of lemongrass (*cymbopogon citratus*) essential oil. *Food Chem Toxicol*, 49, 2268-2272.
50. Bidinotto, L.T., Costa, C.A., Costa, M. et al (2012) Modifying effects of lemongrass essential oil on specific tissue response to the carcinogen N-methyl-N-nitrosourea in female BALB/c mice. *J Med Food*, 15, 161-168.
51. Soares, M.O., Vinha, A., Barreira, S.V., Coutinho, F., Aires-goncalves, S., Oliveira, M.P., Pires, P.C., Castro, A. (2013) Evaluation of antioxidant and antimicrobial properties of the Angolan *cymbopogon citratus* essential oil with a view to its utilization as food biopreservative. *J Agric Sci*, 5(7), 36-45.
52. Balachander, B., Paramasivam, S., Arul kumar, A. (2014) Evaluation of the lemongrass plant (*Cymbopogon citratus*) extracted in different solvents for antioxidant and antibacterial activity against human pathogens. *Asian Pac J Trop Dis*, 4, 134-139.
53. Rathabai, V., Kanimozhi, D. (2013) Phytochemical screening in vitro antioxidant and antimicrobial activity of ethanolic extract of *cymbopogon citratus*. *International Journal of Research in Pharm Med Sci*, 4(3), 760.
54. Cheel, J., Theoduloz, C., Rodriguez, J. et al. (2005) Free radical scavengers and antioxidants from lemongrass (*Cymbopogon citratus* (DC) stapf.). *J Agric Food Chem*, 53(7), 2511-2517.
55. Natchanok, P., Chiravoot, P. (2012) Characterization of eugenol extracted from lemongrass (*Cymbopogon citratus*) for food packaging materials. In: advanced material research (volume 506). Tunkasiri, T. (editor), 603-

606. Doi: 10.4028/www.scientific.net/AMR.506.603
56. Genser, B., Grammer, T.B., Stojakovic, T., Siekmeier, R., Marz, W. (2008) Effect of HMG-CoA reductase inhibitors on low density lipoprotein cholesterol and C-reactive protein: systematic review and meta-analysis. *Int J Clin Pharmacol Ther*, 46, 497-510.
57. Abe, S., Maruyama, N., Hayama, K., Inouye, S., Oshima, F., Yamaguchi, H. 2004. Suppression of neutrophil recruitment in mice by geranium essential oil. *Mediators Inflamm*, 13, 21-54.
58. Gbenou J.D., Ahounou, J.F., Akakpo, H.B., Laleye, A., Yaji, E., Gbaguidi, F., Baba-Moussa, L., Barboux, R., Danson, P., Moudachiron, M., Kotchoni, S.O. (2012) Phytochemical composition of *cymbopogon citratus* and eucalyptus citriodora essential oils and their anti-inflammatory and analgesic properties on Wistar rats. *Mol Biol Rep*, Doi 10.1007/s11033-012-2155-1.
59. Ademuyiwa, A.J., Olamide, O.Y., Oluwatosin, O.O., (2015) The effects of *Cymbopogon citratus* (Lemongrass) on the blood sugar level, lipid profiles and hormonal profiles of Wistar Albino Rats. *Merit Res J Med and Med Sci*, 3(6), 201-216.
60. Hirosh, T., Mitsuy, I., Mi, T., Jin, E.W., Toshiyasu, S., Ikoku, K. (2004) Effect of green tea on blood glucose levels and serum proteomics patterns in diabetic mice and on glucose metabolism in healthy humans. *BMC Pharma*, 4,18-30.
61. Liu, P.X., Pincus, S.M., Takahashi, P.Y., Roebuck, P.D., Iranmanesh, A., Keenan, D.M., Veldhuis, J.D. (2006) Aging attenuates both the regularity and joint syndchroyny of LH and testosterone secretion in normal men: analysis via a model of graded GnRH receptor blockade. *Am J Physiol Endocrinol Metab*, 290(1), E34-E41.
62. Bharti, S.K., Kumar, A., Prakash O, Sharma, N.K., Krishnan, S., Gupta, A.K. (2013) Essential oil of *cymbopogon citratus* against diabetes: Validation by In vivo experiments and computational studies. Open access scientific reports. 2:688 doi:10:4172/scientific reports.688
63. Strasak, A.M., Kelleher, C.C., Brant, L.J., Rapp, K., Ruthmann, E., Concin, H., Diem, G., Pfeiffer, K.P., Ulmer, H., VHM&PP Study Group. (2008) Serum uric acid is an independent predictor for all major forms of cardiovascular death in 28,613 elderly women: a prospective 21-year follow-up study. *Int J Cardiol*, 125, 232-239.
64. Strasak, A.M., Kelleher, C.C., Brant, L.J., Rapp, K., Ruthmann, E., Concin, H., Diem, G., Pfeiffer, K.P., Ulmer, H., VHM&PP Study Group. (2008) Serum uric acid and risk of cardiovascular mortality: a prospective long term study of 83,683 Austrian men. *Clin Chem*, 54, 273-284.
65. Mirghani, M., Liyana, Y., Parveen, J. (2012) Bioactive analysis of lemongrass essential oil. *Int Food Res J*, 19(2), 569-575
66. Omar, B., Mohamed, N., Rahim, Y.A., Wahab, H.A. (2007) Natural flavonoids for the treatment of hyper uricemia, molecular docking studies. *World congress on medical physics and Biochemical engineering IFMBE proceeding volume*, 167-172.
67. Borges, F., Fernande, E., Roleira, F. (2002) Progress towards the discovery of xanthine oxidase inhibitors. *Curr Med Chem*, 9 (2): 195.
68. Ponce, D. A., Glen, J. M. (2002) Relationship of Epithelial Gold Deposits to Large-Scale Fractures in Northern Nevada. *Economic Geology*, 97, 3-9
69. Ravnskov, U. (2000) Hydrocarbon exposure may cause glomerulonephritis and worsen renal failure: evidence based on Hill's criteria of causality. *QJM*, 93:551-550.
70. Anzai, A., Enomoto, A., Endou, H. (2005) Renal urate handling: clinical relevance of recent advances. *Curr Rheumatol Rep*, 7(3): 227-234.
71. Ravnskov, U. (2005) Experimental glomerulonephritis induced by hydrocarbon exposure: A systematic review. *BMC Nephrol*, 6, 15-18
72. Festus, O.O., Dada, F.L., Iweka, F.K., Eyaufe, A.O., Osagie, R.N., Osagie, E.V., et al., (2013) Plasma renal function amongst petrol station attendants in Owerri, South-East Nigeria. *Int J Community Res*, 2; 2515-6562.
73. Awodele, O., Sulayman, A.A., Akintonwe, A. (2014) Evaluation of hematological hepatic and renal function of petroleum tanker drivers in Lagos, Nigeria. *Afri Health Science*, 14; 178-1