

## Original Research Article

## Effect of transfluthrin-impregnated insecticide paper on some biochemical parameters and lung histopathology in rats

I.U. Muhammad<sup>1\*</sup>, A.J. Alhassan<sup>1</sup>, A.A. Imam<sup>1</sup>, A. Idi<sup>1</sup>, A. Mohammed<sup>2</sup>, A. Nasir<sup>3</sup>, A. I. Yaradua<sup>3</sup>, S.M. Adamu<sup>4</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Basic Sciences, Bayero University, P. M. B. 3011, Kano, Nigeria.

<sup>2</sup>Department of Biochemistry, College of Medical Sciences, Abubakar Tafawa Balewa University, P.M.B 0248, Bauchi, Nigeria.

<sup>3</sup>Department of Biochemistry, Faculty of natural and applied Sciences, Umaru Musa Yar'adua University, Katsina, Nigeria.

<sup>4</sup>Department of Pathology, National Orthopaedic Hospital Dala, Kano, Nigeria.

### \*Corresponding Author:

I.U. Muhammad

Email: [ibrahimmuhd@yahoo.com](mailto:ibrahimmuhd@yahoo.com)

**Abstract:** Serum malondialdehyde (MDA), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP) and histological analysis of lungs were carried out in twenty rats exposed to smoke from transfluthrin impregnated insecticide paper. The rats were divided into five groups (I, II, III, IV and V) of four each. Group one served as control while Group II, III, IV and V were exposed to smoke from transfluthrin impregnated paper Insecticide for 15, 30, 60, 120 minutes daily for four weeks respectively. On the 29th day, the rats were euthanized and blood sample was collected and centrifuged for analysis of biochemical parameters (MDA, AST, ALT and ALP), the animals were dissected and lung tissues were collected for histological analysis. Significant increase ( $p < 0.05$ ) in all parameters (MDA, AST, ALT and ALP) were observed in a time dependent pattern compared to normal control. However, histopathological analysis of the lung tissues shows no pathological changes between test groups and normal control. Thus, exposure of rat to smoke from transfluthrin impregnated paper insecticide modifies biochemical parameters (MDA, AST, ALT and ALP) but within the context of duration of this research, no significant pathology was observed in lung tissues. Transfluthrin impregnated paper insecticide should be used with caution.

**Keywords:** Biochemical, histopathological, insecticide, smoke and transfluthrin.

### INTRODUCTION

Malaria is a disease caused by injection of plasmodium parasite into the human body due to bite of certain species of female anopheles mosquitoes. The infective forms (sporozoites) of one or more of at least four different species of plasmodium invade the liver and subsequently red blood cells giving rise to periodic shivering, pyrexia and sweating [1]. Malaria is by far the world's most important tropical parasitic disease killing people more than any other communicable disease except AIDS and Tuberculosis. Worldwide prevalence of the disease is in the order of 350-500 million clinical cases each year, with an estimated annual death of over 1.1 million deaths each year [2]. Mosquitoes are common insect in the family culicidae within the order dipthera, class insecta and phylum arthropoda, there are about 3,500 species of Mosquitoes found throughout the world [3]. In some species of Mosquitoes, the females feed on human and therefore serves as vectors for a number of infectious diseases, human malaria is transmitted only by females of the genus Anopheles, they are approximately 430

Anopheles species out of which only 30-40 transmit malaria (i.e are vectors) in nature. Mosquitoes go through four stages in their life cycle: larva, pupa and adult. Mosquitoes in which the female mouthparts are adapted for piercing and sucking belong to subfamily culicinae, the common house mosquito is classified as culex pipiens, and the species that transmit malaria are classified in the genus anopheles. Insecticides are applied to control larvae (larvicides) or adults (adulticides) mosquitoes. Applications of adulticides or larvicides are made after the presence of mosquitoes has been demonstrated by surveillance procedures. All insecticides must have the name and amount of active ingredient (AI) appearing on the label; examples are DEET and pyrethroids [4]. Pyrethroids are synthetic chemicals similar to natural chemicals pyrethrins produced by the flowers of pyrethrums (*Chrysanthemum cinerariac folium* and *Chrysanthemum cinerariac coccincum*). Pyrethroids now constitute a major proportion of the synthetic insecticides and are common in commercial products such as household insecticide and insect repellent in a

concentration that is generally harmless to human [5]. Insecticide paper is an efficient mosquito killer for indoor use, specially developed to combat malaria by eradication of mosquitoes. The active ingredient in most paper insecticide is the pyrethroid "transfluthrin" which paralyzes the mosquitoes by causing their sodium channels to open in the neuronal membrane. Liver is the largest and heaviest internal organ of the body weighing about 1.4 – 1.6 kg, it is a soft, reddish-brown triangular organ with two lobes, averaging about the size of an American football in adults [6]. Liver performs more than 500 different functions which includes fats, proteins and carbohydrates metabolism. The liver also metabolizes all drugs and other foreign bodies hence, plays a major role in homeostasis [7]. Lungs are paired organs in the chest that perform respiration; it is an essential respiratory organ in all air breathing animals. The lungs are covered by a protective membrane called the pulmonary pleura [8]. The principal function of the lungs is to transport oxygen from the atmosphere into the blood stream and to transport carbondioxide from the bloodstream into the atmosphere [9]. This study assessed the effect of smoke from transfluthrin impregnated insecticide paper on serum malondialdehyde (MDA), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP) and its histopathological effect on rat lungs.

## MATERIALS AND METHODS

### Study animals

Male Albino rats weighing 120 g were obtained from Department of Biological Sciences, Bayero University Kano. Animals were housed in colony cages at an ambient temperature and relative humidity. The animals had free access to standard palletized grower feed and drinking water. The principles of laboratory animal care and guidelines were followed.

### Insecticide paper

The insecticide paper used in the research is a booklet containing sheets of papers treated with transfluthrin 0.45%, and essential oils 2.5% as its active ingredients, each sheet is weighing 0.905g and 16.9cm in length and 7.2cm in width.

## Experimental protocol

This research is an interventional treatment of albino rats with smoke from transfluthrin impregnated insecticide paper. Twenty (20) male albino rats were divided into five groups (I, II, III, IV and V) of four animals each, group one served as normal control while Groups (II, III, IV and V) were exposed to smoke from transfluthrin based impregnated paper Insecticide for 15, 30, 60, 120 minutes daily for 28 days respectively. On the 29<sup>th</sup> day, all the rats were euthanized and blood sample was collected in a dried centrifuge tube for biochemical analysis the carcasses were dissected and from the lungs tissue were collected for histological studies. Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) activity were assayed using Reitman and Frankel method [10], alkaline phosphatase (ALP) activity was assayed using the method developed by Roy [11], serum malondialdehyde was estimated using Ohkawa *et al.* [12] method while the histopathological analysis was done by staining technique according to Auwioro [13].

## Statistical analysis

All data were expressed as mean  $\pm$  standard deviation. Statistical differences between groups were analyzed by one-way analysis of variance (ANOVA) followed by Tukey's test after investigating the data for normality using Shapiro-Wilk test and for variances homogeneity to be sure that the data are normally distributed and variances would be homogenous using GraphPad InStat3 Software version 3.05 Differences of  $P < 0.05$  were considered to be significant [14].

## RESULT


Table 1 present the effect of exposure to graded dosage transfluthrin based insecticide paper on MDA, ALP, ALT and AST concentrations in experimental rats, there is a significant increase in all parameters (MDA, ALP, ALT and AST) in test groups (group II, III, IV and V) compared to normal control (group I) except for ALP and AST which only shows a remarkable increase in groups (III, IV and V). While plate (1-5) shows the histopathological Analysis of lung tissue from experimental rats exposed to graded doses with transfluthrin based paper Insecticide.

**Table-1: Effect of Transfluthrin based Insecticide Paper on MDA, ALP, ALT and AST concentrations in experimental rats.**


Groups	Time of exposure (min)	MDA(nmol/ml)	ALP(IU/L)	AST (IU/L)	ALT(IU/L)
I	-	0.5 $\pm$ 0.17 <sup>a,b,c,d</sup>	43.75 $\pm$ 2.04 <sup>a,b,c</sup>	9.50 $\pm$ 1.30 <sup>a,b,c</sup>	8.50 $\pm$ 2.40 <sup>a,b,c,d</sup>
II	15	0.9 $\pm$ 0.20 <sup>a</sup>	55.78 $\pm$ 2.78	18.25 $\pm$ 2.75	20.28 $\pm$ 4.30 <sup>a</sup>
III	30	1.2 $\pm$ 0.30 <sup>b</sup>	86.20 $\pm$ 3.56 <sup>a</sup>	22.25 $\pm$ 2.08 <sup>a</sup>	23.75 $\pm$ 3.30 <sup>b</sup>
IV	60	1.4 $\pm$ 0.17 <sup>c</sup>	127.00 $\pm$ 3.80 <sup>b</sup>	24.25 $\pm$ 2.22 <sup>b</sup>	24.25 $\pm$ 3.00 <sup>c</sup>
V	120	1.5 $\pm$ 0.13 <sup>d</sup>	126.17 $\pm$ 4.18 <sup>c</sup>	26.50 $\pm$ 3.10 <sup>c</sup>	29.25 $\pm$ 2.60 <sup>d</sup>

Results are presented as Mean  $\pm$  standard deviation, (n=4). Values with the same superscripts in a column are significantly different compared to each other ( $P < 0.05$ )


**Histopathological analysis**




**Plate-1: section of lungs of control rat showing no pathological changes (H and E, mag.  $\times 100$ )**




**Plate-2: section of rat lungs exposed to smoke from transfluthrin based impregnated paper Insecticide for 15 minutes showing no significant pathological changes (H and E, mag.  $\times 100$ )**



**Plate-3: section of rat lungs exposed to smoke from transfluthrin based impregnated paper Insecticide for 30 minutes showing no significant pathological changes (H and E, mag.  $\times 100$ )**



**Plate-4: section of rat lungs exposed to smoke from transfluthrin based impregnated paper Insecticide for 60 minutes showing mild pathological changes (H and E, mag.  $\times 100$ )**



**Plate-5: section of rat lungs exposed to smoke from transfluthrin based impregnated paper Insecticide for 60 minutes showing mild changes (H and E, mag.  $\times 100$ )**

## DISCUSSION

Liver damage is a serious disease characterized by disturbances in the normal functions of the liver. It is clinically diagnosed by determining the serum concentration of liver enzymes (ALT, AST and ALP). ALT and AST are non-plasma specific enzymes involved in transamination of aspartic acid and alanine respectively, the enzymes were reported to reach higher than normal levels in the blood when there is necrosis of the parenchymal cells of the liver as in viral or toxic hepatitis [15]. ALP is also a non-plasma specific enzyme involved in the hydrolysis of a variety of phosphate esters at alkaline PH. These enzymes were reported to reach higher than normal level in the blood in events of impaired liver function [16]. Thus, they are used as serum markers of hepatic damage. Malondialdehyde is used as a measure of lipid peroxidation. Lipid peroxidation is a chain of reaction providing a continuous supply of free radical that initiates further peroxidation [17]. Malondialdehyde is used to determine the indices of oxidative stress in the serum. in this research, significant increase in the level of MDA in transfluthrin impregnated paper insecticide treated groups (II, III, IV and V) was observed compared to the normal control G I this shows that reactive oxygen species were produced when the rats

were expose to doses of transfluthrin impregnated paper insecticide which indicate that there is oxidative stress that depends upon the time of exposure. A significant increase ( $p < 0.05$ ) in serum levels of ALT, AST and ALP was observed in groups (II, III, IV and V) compared to the controls (Group I), this an indication of induction of liver damage caused by the insecticide paper. The enzymes were reported to be higher than normal levels in the blood when there is necrosis of the parenchymal cells of the liver as in viral or toxic hepatitis [18]. This finding support the result of a research carried by Ray [19] who reported that preteroids are generally harmless to human beings in low doses but can harm sensitive individuals. However, no pesticide is 100 percent safe and care must be exercised in the use of any pesticide [20]. The histopathological examination of lungs shows normal cellular architecture with no significant difference between groups. This may be due to the fact that effects in tissue are produce after a prolonged frequent exposure.

## ETHICAL APPROVAL

All authors hereby declare that Principle of laboratory animal care [21] and ethical guidelines for

investigation of experimental pain in conscious animals [22] were observed during experimentation

## CONCLUSION

It can be concluded that within the context of time and dose used in the research, exposure to doses of transfluthrin impregnated paper insecticide lead to liver damage and oxidative stress but does not lead to lungs failure. However, it can be toxic on long time exposure even at lower concentration therefore it should be used with great caution

## REFERENCES

1. World Health Organization. (2015). *Guidelines on the management of latent tuberculosis infection*. World Health Organization.
2. Bruce-Chwatt, L. J., Wernsdorfer, W. H., & McGregor, I. (1988). Malaria: principles and practice of malariology. *Malaria: principles and practice of malariology*.
3. Rueda, L. M. (2008). Global diversity of mosquitoes (Insecta: Diptera: Culicidae) in freshwater. *Hydrobiologia*, 595(1), 477-487.
4. Aranda, C., Eritja, R., & Roiz, D. (2006). First record and establishment of the mosquito *Aedes albopictus* in Spain. *Medical and veterinary entomology*, 20(1), 150-152.
5. Ray, D. E. (1991). Pesticides derived from plants and other organisms. *Handbook of pesticide toxicology*, 2(13), 585-636.
6. Kumar, V., Abbas, A. K., & Aster, J. C. (2013). *Robbins basic pathology*. Elsevier Health Sciences.
7. Guyton AC, Hall JE. Contraction of skeletal muscle. *Textbook of medical physiology*.
9. Heldin, C. H., Landström, M., & Moustakas, A. (2009). Mechanism of TGF- $\beta$  signaling to growth arrest, apoptosis, and epithelial-mesenchymal transition. *Current opinion in cell biology*, 21(2), 166-176.
10. Tarn, D. M., Heritage, J., Paterniti, D. A., Hays, R. D., Kravitz, R. L., & Wenger, N. S. (2006). Physician communication when prescribing new medications. *Archives of Internal Medicine*, 166(17), 1855-1862.
11. Reitman, S., & Frankel, S. (1957). A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *American journal of clinical pathology*, 28(1), 56-63.
12. Roy, A. V. (1970). Rapid method for determining alkaline phosphatase activity in serum with thymolphthalein monophosphate. *Clinical chemistry*, 16(5), 431-436.
13. Ohkawa, H., Ohishi, N., & Yagi, K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical biochemistry*, 95(2), 351-358.
14. Awvioro, O. G. (2002). *Histochemistry and Tissue Pathology Principles and Techniques*. ISBN, 978356279, 154162.
15. Schroeder, S., Enderle, M. D., Baumbach, A., Ossen, R., Herdeg, C., Kuettner, A., & Karsch, K. R. (2000). Influence of vessel size, age and body mass index on the flow-mediated dilatation (FMD%) of the brachial artery. *International journal of cardiology*, 76(2), 219-225.
16. Bruin, A. D. (1976). *Biochemical toxicology of environmental agents*.
17. Adzu, B., Abubakar, M. S., Izebe, K. S., Akumka, D. D., & Gamaniel, K. S. (2005). Effect of *Annona senegalensis* root bark extracts on *Naja nigricotlis nigricotlis* venom in rats. *Journal of ethnopharmacology*, 96(3), 507-513.
18. Santangelo, P. J., Nix, B., Tsourkas, A., & Bao, G. (2004). Dual FRET molecular beacons for mRNA detection in living cells. *Nucleic acids research*, 32(6), e57-e57.
19. Ghosh, J., Das, J., Manna, P., & Sil, P. C. (2010). Acetaminophen induced renal injury via oxidative stress and TNF- $\alpha$  production: therapeutic potential of arjunolic acid. *Toxicology*, 268(1), 8-18.
20. Ray, D. E. (1991). Pesticides derived from plants and other organisms. *Handbook of pesticide toxicology*, 2(13), 585-636.
21. Garrett, W. D., & Barger, W. R. (1979). *U.S. Patent No. 4,160,033*. Washington, DC: U.S. Patent and Trademark Office.
22. NIH. (1996) *Guidelines for the care and use of laboratory animals*. National Academic Press, NIH Publication. 85: 23.
23. Zimmermann, M. (1983). Ethical guidelines for investigations of experimental pain in conscious animals. *Pain*, 16(2), 109-110.