

## The Lipidemic and Antioxidant Role of *Moringa oleifera* Leave Extract Following Salt Loading

Archibong AN, Nku CO, Akwari AA, Esiere KS, Ime AU\*

Department of Physiology, College of Medical Sciences, University of Calabar, Calabar, Nigeria

\*Corresponding author  
Ime AU

### Article History

Received: 08.12.2017

Accepted: 16.12.2017

Published: 30.01.2018

### DOI:

10.21276/sjm.2018.3.1.2



**Abstract:** Salt consumption is essential but has a negative health impact when taken in excess. Therefore, this research is aimed at investigating the possible role of *Moringa oleifera* leave extract on lipids and antioxidant enzymes following salt loading. 24 male albino Wistar rats weighing between 180-240g were used for this study and were divided into four groups (n=6). They were given either normal rat feed and drinking water, high salt diet (8% NaCl diet) + 1% NaCl drinking water and/or *Moringa oleifera* extract (600 mg/kg b.w., orally, once daily). After six weeks of feeding, the animals were sacrificed and blood collected through standard method for analysis. The salt fed untreated rats had significant (p<0.05) increase in Triglycerides (TG) level, Atherogenic indices (P<0.05), Malondialdehyde (MDA), Aspartate aminotransferase (AST) and Alanine aminotransferase (ALT) (P<0.01). These parameters were significantly (p<0.05) reduced to near control values following extract treatment. Salt fed untreated rats were also observed to have significant (P<0.05) reduction in High density lipoprotein (HDL), Catalase (CAT) (P<0.01), Glutathione peroxidase (GPx) (P<0.05), Superoxide dismutase (SOD) and Albumin (ALB) (P<0.01), but the reverse was the case following extract treatment. In conclusion, consumption of *Moringa oleifera* leaf extract ameliorates the adverse effects of salt loading on the atherogenic, antioxidants and other indices thereby preventing possible outcome of cardiovascular disease, atherosclerosis and other related ailment. It is therefore being recommended for further research and possible use in the manufacture of drugs that are necessary in management of cardiac and other related ailments.

**Keywords:** *Moringa oleifera* Lam., Lipid profile, Atherogenic indices, antioxidants, Liver Enzymes.

## INTRODUCTION

*Moringa oleifera* is a tree that is known to have originated from India, it is also being found in Sri-Lanka, Thailand, Pakistan, Phillipine, Indonesia, Taiwan, Haiti, South America, Caribbean and Africa (Nigeria) [10]. It is commonly known as Drumstic, [11] horseradish tree, ben oil (benzoil) [12], mothers best friend and Nebeday depending on location and there are about 13 species of *Moringa* belonging to Moringaceae family [13, 14]. Almost all the plant parts are essential in the synthesis of therapeutic drugs [15], they also have important agricultural, commercial and economical values. Proximate and phytochemical analysis of *Moringa oleifera* leave extract [16] reveals its rich nutritive and bioactive components which include carbohydrate, fat, protein and minerals eg. calcium, iron, magnesium, manganese, phosphorus, potassium, sodium, zinc and water [12, 17, 18]. Others include vitamins (Vit. B1, B2, B3, B5, B6, B9, Vit. C, Vit. E and K), Carotenoids and antioxidants which include flavonoids, glycosides, terpinoids, zeatin [16], quercetin and kaempferol [12, 18]. Antioxidant components of *Moringa oleifera* like Vit. C, carotene and quercetins are known to play

major roles in lowering blood pressure [19, 20, 21], quercetins and flavonoids can inhibit the production of nitric oxide and tumor necrosis factor by Kupffer cell when stimulated by injury [22] flavonoids also protect the cell against injury caused by x-ray and block the progression of the cell cycle and prostaglandin synthesis thereby inhibiting mutation and preventing carcinogenesis in experimental animals [23], the Vit. E antioxidant which is composed of tocopherol and alpha tocopherol is the most abundant and active component of this plant. This vitamins prevent lipid peroxidation chain reaction generated by free radicals from cellular and subcellular membrane which are rich in polyunsaturated lipid thereby preventing atherosclerosis and cancer [24, 25]. The Vitamin C components of *Moringa oleifera* can act as a scavenger of free radicals and do also regenerate Vitamin E indirectly [24], by virtue of this synergy, both vitamins C and E have attracted interest as agent that can retard atherosclerosis by reducing low density lipoproteins oxidation and thus preventing injury to the vascular endothelial cells [26], the antioxidant also protect against structural defect thus inhibiting free radicals formation [39].

Vitamin A is important for normal vision in dim light and for resistance against infection [24, 27, 28] the chlorogenic acid [29], and isothiocyanate component of *Moringa oleifera* also plays a major role in reducing blood sugar level in addition to its anti-inflammatory [26, 30, 31] anticancer and antimicrobial effect [32-34]. *Moringa oleifera* extract are also known to possess antitumor and hepatoprotective activities [30, 35], antispasmodic [14, 36, 37] and antiepileptic activities [38]. In addition, blood parameters like Red blood cell (RBC) count, packed cell volume (PCV), hemoglobin (Hb), white blood cell (WBC) counts, and platelets (Plt) were shown to be enhanced following consumption of *M. oleifera* [40] and even in situations of excessive salt consumption [41]. This research is therefore aim at investigating the possible role of *Moringa oleifera* leave extract on lipids and antioxidant enzyme following salt loading. This research was pre-conceive taking full cognizance of the fact that excessive dietary salt consumption over an extended period of time in our day to day life may predispose an individual to hypertension, cardiovascular disease, left ventricular hypertrophy and other opportunistic diseases, [1-7] it has also been shown to increase the risk of myocardial infarction, stroke, arterial stiffness and heart failure [8-9].

## MATERIALS AND METHODS

### Experimental-Animals

Twenty four (24) male albino Wistar rats weighing initially between 180 to 250g obtained from the animal house of the Department of Physiology, University of Calabar, Nigeria were employed for this study for 6 weeks. The animals were allowed free access to their feed and drinking water. The rats were weighed before commencement of the feeding experiment and thereafter were weighed daily. Ethical approval was obtained from the Ethics Committee of the Faculty of Basic Medical Sciences, University of Calabar, Nigeria. They were nursed under control and the environmental conditions were in accordance with international standard [42].

### *Moringa oleifera* extract preparation

The aqueous extract was prepared according to standard procedure [43]. Fresh leaves of *Moringa oleifera* were obtained from Calabar municipality, Cross River State and were identified by the Herbarium in Botany Department. The leaves were washed to remove debris and were later dried in an airy-room away from direct sunlight to avoid possible damage to their phyto-constituents for two days. The leaves were further oven-dried for 30 minutes at the temperature of 40°C. The dried leaves were grinded to powdered form.

About 1400gram of the powdered *Moringa oleifera* leaf was soaked in 7000ml of distilled water for about 24 hours. The mixture was then filtered with a white cotton (satin) material, followed with filter paper (Whatmann No.1) into beakers and placed in an oven.

The filtrate was evaporated to dryness using a rotary evaporator with temperature set at 50°C. *Moringa oleifera* extract was then collected into a sample bottle and preserved in a refrigerator.

### Salt diet and Drinking water

Salt feed containing 8% NaCl was prepared by mixing together 8g NaCl in 92g of the rat feed. Also 1% NaCl drinking water was prepared by dissolving 100g of NaCl in small quantity of distilled water and volume made upto 10L with distilled water [45].

### Experimental design

Twenty four (24) male albino Wistar rats weighing between 180 – 240g was randomly assigned into four (4) groups of six (6) rats each.

Group 1 (control) - received normal rat feed + drinking water

Group 2- received same as group 1 + *Moringa oleifera* extract (600mg/kg o.p. once daily)

Group 3- received 8% NaCl diet + 1% NaCl water

Group 4- received same as group 3 + *Moringa oleifera* extract (600mg/kg o.p. once daily).

The administration was done orally and the experiment lasted for a period of six weeks.

### Collection of blood samples and analysis of different parameters

Blood samples were collected via cardiac puncture into EDTA capped bottles and the different parameters were analyzed thus.

### Measurement of Lipid Profile

#### Measurement of total cholesterol

Total cholesterol was measured according to standard procedure [44].

#### Principle

Cholesterol esters are hydrolysed by cholesterol esterase to produce cholesterol and fatty acids. The cholesterol is oxidized by cholesterol oxidase to cholesterolone and hydrogen peroxide. The H<sub>2</sub>O<sub>2</sub> is later hydrolysed by peroxidase to form water and oxygen. The oxygen then reacts with 4-aminoantipyrine which is the chromogen to form quinoneimine. The colour intensity of the solution is proportional to the concentration of cholesterol in the sample.

The samples were mixed and incubated for 10 minutes in a water bath at 37°C. The colour produced was read colorimetrically at 540nm.

#### Calculation

$$\frac{\text{Absorbance of test X concentration of standard}}{\text{Absorbance of standard}}$$

### **Measurement of triglyceride**

Triglyceride was measured according to standard procedure [44].

#### **Principle**

Triglyceride in the sample was hydrolysed by lipoprotein lipase to glycerol and frees fatty acids. Glycerol is phosphorylated by the kinase to form glycerol-3-phosphate and ATP. The glycerol phosphate is then oxidized by glycerolphosphate oxidase to dihydroxyacetone phosphate and H<sub>2</sub>O<sub>2</sub>. The H<sub>2</sub>O<sub>2</sub> is hydrolysed by peroxidase to form H<sub>2</sub>O and O<sub>2</sub>. The O<sub>2</sub> then react with 4-amino-antipyrine and phenol to form the colour complex quinoneimine.

The samples were mixed and incubated for 10 minutes in a water bath at 37°C. The colour produced was read colorimetrically at 540nm.

#### **Calculation**

$$\frac{\text{Absorbance of test X Concentration of standard}}{\text{Absorbance of standard}}$$

### **Measurement of high density lipoprotein cholesterol**

The High Density Lipoprotein was measured according to standard procedure [44].

#### **Principle**

HDL-cholesterol is a precipitate of apo protein B-containing lipoprotein using a mixture of sodium phosphotungstic acid and magnesium chloride.

The samples were mixed thoroughly and allowed to stand at room temperature for 15 minutes and later centrifuged at 3000 revolutions per minute.

The samples were mixed and incubated for 10 minutes in a water bath at 37°C.

#### **Calculations**

$$\frac{\text{Absorbance of test X concentration of standard}}{\text{Absorbance of standard}}$$

Final result is multiplied by the dilution factor 3.0

### **Measurement of low and very low density lipoprotein**

Low density and Very low density lipoprotein concentrations were measured based on Friedwald formular [46].

$$\text{VLDL}_C = \frac{\text{Triglyceride}}{2.22}$$

$$\text{LDL}_C = \text{Total Cholesterol} - (\text{HDL}_C + \text{VLDL}_C)$$

### **Measurement of Cardiac risk ratio (Atherogenic indices):**

The different cardiac risk ratios were measured according to standard procedure.

$$\text{Cardiac Risk Ratio (CRR)} = \text{Tc}/\text{HDL-c} \quad [47]$$

$$\text{Atherogenic coefficient (AC)} = (\text{Tc} - \text{HDL-c}) / \text{HDL-c} \quad [48]$$

$$\text{Atherogenic index of plasma (AIP)} = \log (\text{TG} / \text{HDL-c}) \quad [49]$$

### **Measurement of Antioxidants**

The different antioxidants were analyzed according to standard procedure. Glutathione Peroxidase (GPx) activity was determined using the Rice Evans method [50] Malondihaldihyde (MDA) was determined according to standard procedure [51]. Catalase (CAT) activity in the serum was determined as described by Abebi [52]. The change in the absorbance was monitored spectrophotometrically at 240nm over a 5min period.

Superoxide Dismutase (SOD) activity in the serum was determined by assessing the inhibition of pyrogallol-auto oxidation [53]. Changes in the absorbance at 40nm were recorded at 1min interval for 5min. SOD activity was determined from standard curve of % inhibition of pyrogallol auto-oxidation with SOD activity

### **Measurement of liver enzymes**

#### **Measurement of Aspartate aminotransferase (AST) and Alanine aminotransferase (ALT)**

AST and ALT were measured as described by Reitman and Frankel [54] and as used by Archibong *et al.* [55].

#### **Principle**

AST hydrolysed aspartic acid and  $\alpha$ -ketoglutarate substrate to give oxaloacetic acid and glutamate. The oxaloacetic acid is spontaneously decarboxylated to form pyruvate which reacts with 2,4-dinitrophenylhydrazine (DPNH) to give a brown-coloured hydrozone, which is read in the colorimeter at 540nm [54].

ALT reacts with alaninic acid and  $\alpha$ -ketoglutarate to yield pyruvate and glutamate. The pyruvate formed reacts with 2, 4-dinitrophenylhydrazine (DPNH) to give a brown-coloured hydrazone, which is read in the colorimeter, at 540nm [54].

Each test tube containing the sample was incubated in a water bath for 30 minutes. The test-tubes were then removed and kept on a working bench. 0.5ml of 2,4 - nitrophenylhydrazine was added to each test tube, mixed thoroughly and allowed on the bench for 20 minutes. 0.4N NaOH was added to stop the reaction which was read colometrically at 540nm.

## Calculations

$$\frac{\text{Absorbance of test} \times \text{Concentration of AST standard}}{\text{Absorbance of standard}}$$

## Measurement of alkaline phosphatase (ALP)

ALP was measured using standard method [83].

## Principle

P- nitrophenyl phosphate is hydrolysed by alkaline phosphate to give phosphate and p-nitrophenol. The p-nitrophenol complex formed is read colometrically at 405nm.

The samples were mixed thoroughly and absorbance read immediately.

## Calculations

$$\frac{\text{Absorbance of test} \times \text{Concentration of standard}}{\text{Absorbance of standard}}$$

## Measurement of Serum Proteins

### Total protein

The total protein was measured using the Burette method as described by Tiez [56]

### Principle

The peptide bond in protein has affinity for burette reagent. In alkaline medium,  $\text{CuSO}_4$  in the burette reagent reacts with the peptide bond of protein to give a blue colour complex. The colour produce is proportional to the concentration of protein in the sample. The samples were thoroughly mixed and incubated in a water bath at  $37^\circ\text{C}$  for 10 minutes. After which they were then removed from the water bath and read colometrically at 540nm [56].

### Serum albumin

The serum albumin was measured using the Bromocresol green method as described by Grant [57]

### Principle

In an acidic medium bromocresol green dye binds to albumin to give a violet green colouration. Samples were mixed thoroughly and incubated at room temperature for 5 minutes and read colometrically at 620nm [57].

### Globulin

Calculation: Total Protein – Albumin = Globulin

## STATISTICAL ANALYSIS

Data are presented as mean  $\pm$  SEM. Data were analyzed using one-way analysis of variance (ANOVA) and then followed by post hoc test (least square deviation). Data analysis was done with the help of computer software (Excel and SPSS version 17.0 for

windows). P-values of less than 0.05 were considered as significant.

## RESULTS AND DISCUSSIONS

### Lipid profile

As shown in Table 1 the difference in total cholesterol concentration in the *M. oleifera* group ( $181.58 \pm 4.88$ ), Salt fed group ( $206.04 \pm 12.21$ ) and Salt fed + *M. oleifera* group ( $222.58 \pm 24.04$ ) was of no statistical significance when compared with the control group ( $208.11 \pm 8.81$ ) respectively. Triglyceride concentration in the salt fed groups ( $27.61 \pm 0.46$ ) was significantly increased ( $P < 0.05$ ) when compared with the control ( $23.25 \pm 1.74$ ) and *M. oleifera* ( $24.95 \pm 1.13$ .) groups respectively but treatment with the extract reduced the TG concentration in salt fed group as indicated by its concentration in the salt fed + *M. oleifera* group ( $25.73 \pm 1.02$ ).

High density lipoprotein (HDL) concentration in the salt fed group ( $17.53 \pm 1.09$ ) was significantly decreased ( $p < 0.05$  and  $p < 0.001$ ) when compared with that of control ( $21.85 \pm 1.01$ ) and *M. oleifera* ( $24.53 \pm 1.95$ ) groups respectively but treatment with the extract increased the concentration of HDL concentration in the salt fed group as indicated by its concentration in the salt fed + *M. oleifera* group ( $21.05 \pm 0.54$ ).

The difference in Low density lipoprotein (LDL-c) concentration in the *M. oleifera* groups ( $151.14 \pm 4.69$ ), salt group ( $182.99 \pm 12.12$ ) and Salt fed + *M. oleifera* ( $196.39 \pm 24.25$ ) was of no statistical significance when compared with the control group ( $181.68 \pm 9.20$ ).

The very low density lipoprotein concentration (VLDL-c) in the *M. oleifera* group ( $5.91 \pm 0.61$ ) was significantly higher ( $P < 0.05$ ) than that of the control group ( $4.58 \pm 0.37$ ), salt fed group ( $5.52 \pm 0.09$ ) and Salt fed + *M. oleifera* group ( $5.15 \pm 0.20$ ) respectively

### Cardiac risk indices

As shown in Table 2, the cardiac risk ratio in the salt fed group ( $11.93 \pm 0.96$ ) was significantly increased ( $p < 0.05$  and  $p < 0.01$ ) when compared with the control ( $9.71 \pm 0.84$ ) and *M. oleifera* ( $7.62 \pm 0.58$ ) groups respectively but treatment with the extract decreased the CRR in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group ( $10.63 \pm 1.21$ )

The atherogenic coefficient in the salt fed group ( $10.93 \pm 0.96$ ) was significantly increased ( $p < 0.05$  and  $p < 0.01$ ) when compared with the control ( $8.71 \pm 0.84$ ) and *M. oleifera* ( $6.62 \pm 0.58$ ) groups respectively but treatment with the extract decreased the atherogenic coefficient in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group ( $9.63 \pm 1.21$ )

The atherogenic index in the salt fed group (0.20±0.02) was significantly increased ( $p<0.05$  and  $p<0.001$ ) when compared with the control (0.02±0.05) and *M. oleifera* (0.01±0.04) groups respectively but treatment with the extract decreased the atherogenic index in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (0.09±0.02).

#### Antioxidant Enzymes

As shown in Table 3, catalase (CAT) level in the salt fed group (0.27±0.03) was significantly decreased ( $p<0.01$  and  $p<0.001$ ) when compared with control (0.37±0.01) and *M. oleifera* (0.41±0.02) groups respectively but treatment with the extract increased the Catalase level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (0.30±0.03).

The Glutathione Peroxidase (GPx) level in the salt fed (850.67±42.33) group was significantly decreased ( $p<0.01$ ) when compared with control (946.69±31.17) and *M. oleifera* (1057.67±53.36) groups respectively but treatment with the extract increased the GPx level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (888.17±36.84)

The Superoxide dismutase (SOD) level in the salt fed (0.12±0.02) group was significantly decreased ( $p<0.001$ ) when compared with control (0.22±0.02) and *M. oleifera* (0.23±0.02) groups respectively but treatment with the extract increased the SOD level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (0.16±0.01).

The malondialdehyde (MDA) level in the salt fed (15.33±0.84) group was significantly increased ( $p<0.01$ ) when compared with control (12.00±0.77) and *M. oleifera* (11.67±0.49) groups respectively but treatment with the extract decreased the MDA level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (13.80±0.83).

#### Liver Enzymes

As shown in Table 4 the Aspartate Transferase (AST) level in the salt fed group (59.28±4.10) was significantly increased ( $p<0.01$  and  $p<0.001$ ) when compared with that of control (43.16±2.08) and *M.*

*oleifera* (46.25±0.25) groups respectively but treatment with the extract decreased the AST level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (47.69±2.31).

Alanine Transferase (ALT) level in the salt fed group (70.33±3.91) was significantly increased ( $p<0.001$ ) when compared with that of control (47.28±3.26) and *M. oleifera* (45.54±3.82) groups respectively but treatment with the extract decreased the ALT level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (49.13±2.10)

Alkaline Phosphatase (ALP) level in the salt fed group (44.76±3.17) was significantly decreased ( $p<0.01$ ) when compared with that of control (47.41±2.51) but treatment with the extract further decreased the ALP level in the *M. oleifera* (35.69±3.06) and salt fed + *M. oleifera* group (42.51±0.83).

#### Serum Protein

As shown in Table 5, the total protein concentration in the salt fed group (60.17±0.80) was significantly decreased ( $p<0.05$  and  $p<0.01$ ) when compared with that of control (64.33±0.80) and *M. oleifera* (64.17±0.40) groups respectively but treatment with the extract decreased the total protein level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (62.00±0.73).

The Albumin level in the salt fed group (26.33±0.61) was significantly decreased ( $p<0.001$ ) compared with that of *M. oleifera* (34.00±0.73) and control group (34.33±0.61) but treatment with the extract increased the albumin level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (30.00±0.73).

Globulin level in the salt fed group (34.17±0.17) was significantly increased ( $p<0.001$ ) when compared with that of control (30.00±0.42) and *M. oleifera* (30.17±0.49) groups respectively but treatment with the extract decreased the globulin level in the salt fed group as indicated by its value in the salt fed + *M. oleifera* group (32.50±0.62).

**Table-1: Showing comparison of Lipid profile in the different groups**

	Tc	TG	HDLc	VLDLc	LDL
Control	208.11±8.81	23.25±1.74	21.85±1.01	4.58±0.37	181.68±9.20
<i>M. oleifera</i>	181.58±4.88	24.95±1.13	24.53±1.95	5.91±0.61*	151.14±4.69
Salt Fed	206.04±12.21	27.61±0.46*	17.53±1.09*	5.52±0.09	182.99±12.12
Salt Fed + <i>M. oleifera</i>	222.58±24.04	25.73±1.02	21.05±0.54	5.15±0.20	196.39±24.25

Values are represented as Mean ± SEM. \* $p<0.05$  vs Control.

**Table-2: Showing comparison of Atherogenic indices in the different groups**

	CRR	AC	AIP
Control	9.71±0.84	8.71±0.84	0.02±0.05
<i>M. oleifera</i>	7.62±0.58	6.62±0.58	0.01±0.04
Salt Fed	11.93±0.96	10.93±0.96	0.20±0.02*
Salt Fed + <i>M. oleifera</i>	10.63±1.21	9.63±1.21	0.09±0.02*

Values are represented as Mean ± SEM. \*p<0.05 vs control.

**Table-3: Showing comparison of Antioxidant enzymes in the different groups**

	MDA	CAT	GPx	SOD
Control	12.00±0.77	0.37±0.01	946.69±31.17	0.22±0.02
<i>M. oleifera</i>	11.67±0.49	0.41±0.02	1057.67±53.36	0.23±0.02
Salt Fed	15.33±0.84**	0.27±0.03**	850.67±42.33	0.12±0.02***
<i>M. oleifera</i> + Salt fed	13.83±0.83	0.30±0.03*	888.17±36.84	0.16±0.01*

Values are represented as Mean ± SEM. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 vs control.

**Table-4: Showing comparison of liver enzyme in the different groups**

	AST	ALT	ALP
Control	43.16±2.08	47.28±3.26	47.41±2.51
<i>M. oleifera</i>	43.25±0.25	45.54±3.82	35.69±3.06
Salt Fed	59.28±4.10***	70.33±3.91***	44.76±3.17***
Salt fed + <i>M. oleifera</i>	47.69±2.31	49.13±2.10	42.51±0.83

Values are represented as Mean ± SEM. \*\*\*p<0.001 vs control

**Table-5: Showing comparison of Plasma Protein in the different groups**

	Total Protein	Albumin	Globulin
Control	64.33±0.80	34.33±0.61	30.00±0.42
<i>M. oleifera</i>	64.17±0.40	34.00±0.73	30.17±0.49*
Salt Fed	60.17±0.40**	26.00±0.61	34.17±0.17***
Salt Fed + <i>M. Oleifera</i>	62.00±0.73***	30.00±0.73***	32.50±0.62***

Values are represented as Mean ± SEM. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 vs control

The result as obtained from this investigation is quite amazing and shows to a large extent the potent effect of *M. oleifera* extract.

The lipid profile analysis revealed that extract of *M. oleifera* was able to significantly reverse the increase in TG and the decrease in HDL-c evidenced in salt loaded rats. This result is further corroborated by earlier researches by Mehta *et al.*, [84, 85] which shows that extract of *M. oleifera* was able to bring about increase in HDL but decrease in TG, vLDL, LDL and atherogenic indices. Raised levels of serum total cholesterol, triglycerides and low density lipoprotein cholesterol are possible indication of coronary heart attack, risk of heart disease and stroke [58]. This result is of significance because accumulation of LDL-c predisposes one to cardiovascular disease, but on the contrary accumulation of HDL-c prevents against cardiovascular disease. The ability of *M. oleifera* leaf extract to reduce these bad cholesterol in the blood maybe unconnected with the fact that leaf extract of *M. oleifera* contain omega-3 and omega-6 fatty acid [59] which is believed to mediate the decrease in the concentrations of bad cholesterol thereby preventing atherosclerosis and related cardiovascular diseases. The increase in HDL-c observed in the *M. oleifera* treated

groups could also be attributed to omega 3 component of the extract [60, 61], which is equally important because HDL-c is the good cholesterol that function in preventing the accumulation of bad cholesterol and ameliorating the risk of heart disease. It has also been shown that low concentration of HDL increases the risk of atherosclerotic disease therefore individuals with seemingly low level of LDL also stand the same risk in situations of inadequate HDL [62 - 64].

It is also important to note that alpha tocopherol which is the most abundant active component of this plant functions effectively in preventing lipid peroxidation chain reaction generated by free radicals from cellular and subcellular membrane which are rich in polyunsaturated lipid thereby preventing atherosclerosis and cancer [24, 25]. The Vitamin C components of *Moringa oleifera* can act as a scavenger of free radicals and do also regenerate Vitamin E indirectly [24], by virtue of this synergy, both vitamins C and E have attracted interest as agent that can retard atherosclerosis by reducing low density lipoproteins oxidation and thus preventing injury to the vascular endothelial cells [26], Therefore the ability of *M. oleifera* leaf extract to reduce these bad cholesterol following salt loading

shows that their consumption would be beneficial to health.

Atherogenic index is a strong marker that predicts the risk of atherosclerosis and coronary heart disease [89], it reflects the true relationship between protective and atherogenic lipoprotein and is associated with the size of pre and anti atherogenic lipoprotein particles [90]. A decrease in atherogenic index as shown by the *M. oleifera* extract treated group shows its potent health effect. This is because a lower value for atherogenic indices is associated with lower risk of cardiovascular diseases but a higher value indicate higher risk [90, 91] This is further being supported by earlier research study by Emmanuel *et al.* [84, 86] which shows that extract of *M. oleifera* reduces elevation in cardiac risk ratio. This point goes a long way to explain the reason why administration of the extract brought about an increase in HDL.

Following the lipid profile result we decided to further investigate the effect of this extract on antioxidant enzymes following salt loading. The antioxidant result revealed that there was a significant decrease in catalase and superoxide dismutase activities caused by salt loading. But this was significantly reversed following combine treatment with *M. oleifera* extract. Also the abnormal increase in Malondialdehyde caused by salt loading was significantly reversed following the administration of *M. Oleifera* extract. This result is consistent with earlier findings by Bonoy *et al.* [65] which shows that extract of *M. oleifera* decreased MDA concentration while bringing about increase in CAT, SOD and GPx concentrations when administered alongside with high salt in rats [87, 88]. MDA is an important Marker for oxidative stress and lipid peroxidation [66, 67], that its activities were significantly increased in the salt fed group reveals to what extent there was peroxidation also ROS and free radicals may have been generated showing how detrimental excessive salt intake could be to the system.

CAT, SOD and GPx are important antioxidant enzymes that aids in the scouting and degrading of free radicals. SOD perform important function by catalyzing the conversion of superoxide radicals to O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> [68, 69] while Catalase further decomposes the toxic H<sub>2</sub>O<sub>2</sub> to O<sub>2</sub> and H<sub>2</sub>O [70]. GPx is also known to catalyze the reduction of hydroperoxide by glutathione [71], it also protects against oxidative stress. That the level of antioxidant enzymes was reduced following Salt loading maybe unconnected with the fact that most of the antioxidant enzymes must have been used up in the degrading of the free radicals that were hitherto generated as a result of excessive salt ingestion or their production may have been inhibited following salt loading. Consequently the antioxidant enzyme activities were shown to have appreciated immensely following the administration of *M. oleifera* extract, as evident by the results. This edible plant leaves have been shown to

possess important antioxidant components [13, 72] eg flavonoids, terpenoids, [16], quercetin and kaempferol [12, 18] which are believed to have boosted the activities of the antioxidant enzymes [73] as experienced by the *M. oleifera* + Salt treated group. Quercetins and flavonoids can inhibit the production of nitric oxide and tumor necrosis factor by Kupffer cell when stimulated by injury [22] flavonoids also protect the cell against injury caused by x-ray and block the progression of the cell cycle and prostaglandin synthesis thereby inhibiting mutation and preventing carcinogenesis in experimental animals [23], The Vitamin C and E components of *Moringa oleifera* can act as a scavenger of free radicals [24]. Therefore this result goes a long way to proof how usefull *M. oleifera* can be in maintaining good health.

The serum enzyme results revealed that the hepatocytes or liver tissues benefited positively from the *M. oleifera* extract administration. This is because the increase in ALT and AST caused by ingestion of excess salt as revealed by the result was significantly reduced following the administration of *M. oleifera* extract. This result is further corroborated by earlier researches published by Ezejindu *et al.* and Afzal *et al.* [77,92], which shows that *M. oleifera* has a protective effect against liver injury. It is important to note that AST, ALT and ALP are biomarkers for healthy liver [74] and they play major role in the diagnosis of heart and liver disease including infections, [75] therefore any abnormal increase in AST, ALT and ALP is suggestive of tissue damage [76]. That this extract of *M. oleifera* was able to reverse the increase in liver enzymes occasion by salt loading, point to the ability of the extract to prevent hepatic damage as reported by Ezejindu *et al.*[77]. Liver protective herbal drugs contain a variety of chemical constituents like phenols, coumarins, lignans, essential oil, monoterpenes, carotinoids, glycosides, flavonoids, organic acids, lipids, alkaloids and xanthenes [93], most of these are present in the *Moringa oleifera* [17] leaves extract and so may be responsible for this beneficial effect. We can infer that leave extract of *Moringa oleifera* has an appreciable ability to prevent damage to the liver, therefore its ingestion may be of immense benefit to health in general.

The serum protein result shows that *M. oleifera* leaf extract was able to significantly reverse the decrease in total protein occasion by excessive ingestion of salt. This result is in consistence with findings by Dharmendra *et al.*, [78] which revealed that administration of *M. oleifera* was able to increase total protein level. This increase in total protein concentration may be due to enhanced protein synthesis in the hepatic cells arising from inhibition of lipid peroxidation [79]. The globulin level was decreased following extract administration. This is of significant because elevated globulin concentration in plasma is an indication of stimulation of the immune system [80,

81]. The serum albumin result reveals that no destruction on the hepatocytes [82] was observed following moringa extract administration but the reverse was the case following salt loading, this result is very essential since albumin levels has been shown to be lowered in chronic and acute liver disease by Burtis *et al.*, [81]. Therefore that decrease in albumin level following salt loading was reversed following moringa extract administration is an indication that this extract may possess certain components that may prevent the liver from being destroyed by excessive salt ingestion.

## CONCLUSION

In conclusion consumption of Moringa oleifera leaf extract ameliorates the adverse effects of salt loading on the atherogenic, antioxidant enzymes and other indices thereby preventing possible outcome of cardiovascular disease, atherosclerosis and other related ailment. It is therefore being recommended for further research and possible use in the manufacture of drugs that are necessary in management of cardiac and other related ailments.

## ACKNOWLEDGEMENTS

The authors of this article do sincerely appreciate the effort of all those who supported this research in different ways. We want to say a big thank you to Mr. Ededet Umoh of Physiology Department who helped to supply the rats, breed and made them available for sacrifice and also to Mrs. Irene Bassey who made all the reagents and equipment available for use during the course of the study. Finally we want to thank the head of department of Physiology for allowing us to use the laboratory and other facilities for the study

## AUTHORS CONTRIBUTION

This work was carried out in collaboration between all authors. Author ANA wrote the first draft of the manuscript, managed the literature and performed the statistical analysis, author CON and AUI designed the study, wrote the protocol and edited the manuscript, while authors AAA and KSE contributed in carrying out the feeding regimens and analysis of blood samples. All authors read and approved the final manuscript.

## REFERENCES

1. Blaustein, M. P. (1977). Sodium ions, calcium ions, blood pressure regulation, and hypertension: a reassessment and a hypothesis. *The American journal of physiology*, 232(5), C165–73.
2. Sacks, F. M., Svetkey, L. P., Vollmer, W. M., Appel, L. J., Bray, G. A. Harsha, D., Obarzanek, E., Conlin, P. R., & Miller, E. R. (2001). Effects on Blood Pressure of Reduced Dietary Sodium and the Dietary Approaches to Stop Hypertension (DASH) Diet. *New England Journal of Medicine*, 344 (1), 3–10.
3. Karanja, N., Erlinger, T. P., Pao-Hwa, L., Miller, E. R., & Bray, G. A. (2004). The DASH diet for high blood pressure: from clinical trial to dinner table. *Cleveland Clinic Journal of Medicine*, 71 (9), 745–53.
4. Cappuccio, F. P. (2007) Salt and cardiovascular disease. *BMJ*, 334 (7599), 859–60.
5. Strazzullo, P., D'Elia, L., Kandala, N. B., & Cappuccio, F. P. (2009). Salt intake, stroke, and cardiovascular disease: meta-analysis of prospective studies. *BMJ*, 339, b4567.
6. Mugavero, K. L., Gunn, J. P., Dunet, D. O., & Bowman, B. A. (2014). Sodium Reduction: An Important Public Health Strategy for Heart Health. *Journal of Public Health Management Practice*, 20 (101), S1–S5.
7. Barbara, E. M., Steve, A., Lucile, A. C., Cheryl, A. A., Thomas, B. J., Wayne, W. C., Steven, C., Frank, H., Miriam, N., Marian, L. N., Rafael, P. E., Anna, M. S., Mary, S., & Alice, H. L. (2016). The 2015 Dietary Guidelines Advisory Committee Scientific Report: Development and Major Conclusions". *Advanced Nutrition*. 7, 438–444.
8. Susic, D; Frohlich, E. D. (2012). Salt consumption and cardiovascular, renal, and hypertensive diseases: clinical and mechanistic aspects. *Current opinion in lipidology*, 23 (1), 11–6.
9. He, F. J., & MacGregor, G. A. (2010) "Reducing Population Salt Intake Worldwide: From Evidence to Implementation". *Progress in Cardiovascular Diseases*, 52 (5), 363–382.
10. Ted, R., & Elevitch C.R. (2011). Farm and forestry production and marketing profile for moringa. In: Specialty crop for pacific. Island Agroforestry Holualoa, Hawaii Permanent Agriculture Resources.
11. Martins, J. V., Marshak, A., Remer, L., Rosenfeld, D., Kaufman, Y. J., & Fernandez, B. R. (2007). Remote sensing the vertical profile of cloud droplet effective radius, thermodynamic phase and temperature. *Amos Chem Phys Disc.*, 7, 4481-4519.
12. Leone, A., Spada, A., Battezzati, A., Schiraldi, A., Aristil, J., & Bertoli, S. (2005). Cultivation genetic ethnopharmacology of Moringa oleifera leaves. An overview. *International Journal of Molecular Science*, 16(6), 12791-12835.
13. Anwar, F., Latif, S., Ashraf, M., & Gilani, A. (2007). Moringa oleifera: a food plant with multiple medicinal uses. *Phytotherapy Research*, 21(1), 17-25.
14. Patel, S., Thakur, A., Chanday, A. (2010). Review of three medicinal and economical importances to the health and nationday. *Drug Invent Today*, 2(7), 339-342.
15. Wadhwa, S., Panwar, S., Saini, N., Rawat, S., & Singhal, M. (2013). A review on commercial traditional uses phytoconstituents and pharmacological activity of *Moringa oleifera*. *Global Journal of Traditional Medicine*, 2, 1-13.
16. Fahey, J. (2005). A review of the medical evidence for its nutritional, therapeutic and prophylactic properties. *Trees Life J.*, 2(4), 1-5.

17. Vinoth, B., Manivasagaperumal, R., & Balamurugun, S. (2012). Phytochemical analysis and antibacterial activity of *Moringa oleifera* Lam. *Int J Res Biol Sci.*, 2(3), 98- 102
18. Gopalan, C., Rama, B., & Balasubramanian, V. (1989). Nutritive value of Indian food. National Institute of Nutrition Indian Council of Medical Research.
19. Newton, K. A., Richard, N. B., & Gladys, M. T. (2010). Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* Lam grown in Ghana. *Food Chemist*, 122(4), 1047-1054.
20. Julia, P. C., Yangping, X., Hong, C., Min-Hsiung, P., Chi-King, H., & Rodolfo, J. (2013). Determination of flavonoid by LC/MS and anti-inflammatory activity of *Moringa oleifera*. *J Funct Food*, 5(4), 1892-1899.
21. Larson, A. J., Symon, J. D., & Jalili, T. (2012). Therapeutic potential of quercetin to decrease blood pressure: review of efficacy and mechanism. *Adv Nutr.*, 3(1), 39-46.
22. Kawada, N., Seki, S., Inove, M., & Kuroki, T. (1998). Effect of antioxidants quercetin and N-acetylcystein on the function of cultured rats hepatic cell stellate cell and kupffer cells. *Hepatology*, 27(5), 1265-1274.
23. Abdulla, M., & Gruber, P. (2000). Role of diet modification in cancer prevention. *Biofactor*, 12(1-4), 45-51.
24. Kummur, V., Abbas, A. K., Fausto, N., Robbsins, S. L., & Contran R. S. (2004.). Pathology bases of disease. New York. WB Saunder
25. Nordberg, J., & Ainer, A. (2001). Reactive oxygen species antioxidants and the mammalian thioredoxin. *Free Radical Bio Med.*, 31(11), 1287-1321.
26. Cheenpracha, S., Park, E. J., Yoshida, W. Y., Barit, C., Wall, M., Pezzuto, J., & Chang, L. C. (2010). Potential anti-inflammatory phenolic glycoside from the medicinal plant *Moringa oleifera* fruit. *Bioorg Med Chem.*, 18(17), 6598-6602.
27. Sudha, P. S., Basheeruddin, A., Sunil S. D., & Gowda K. C. (2010). Immunomodulatory activity of methanolic leaf extract of *M. oleifera* in animals' institute of animal health and veterinary biologicals, Hebbal, Bangalore. *Ind J Physiol Pharmacology*, 54(2), 133-140.
28. Stephensen, C. B. (2001). Vit A, infection and immune function. *Ann Rev Nutr.*, 21, 167-192.
29. Tunnicliffe, J. M., Eller, L. K., Reimer, R. A., Hettel, D. S. & Shearer, J. (2011). Chlorogenic acid differentially affects postparandial glucose and glucose dependent insulinotropic polypeptides response in rats. *Appl Physiol Nutr Metab.*, 36(5), 650-659.
30. Sharma, R., & Vaghela, J. (2011). Anti-inflammatory activity of *Moringa oleifera* leaf and pod extracts against carrageenen induced paw edema in albino mice. *Pharmacol Online*, 1, 140-144.
31. Mahajan, S. G., & Mehta, A. A. (2010). Immunosuppressive activity of ethanolic extract of seed of *Moringa oleifera* Lam in experimental immune inflammation. *J Ethnopharmacology*, 130(1), 183-186
32. Moon, K., Guallar, E., & Navas-Acien, A. (2012). Arsenic exposure and cardiovascular disease. An updated systemic review. *Curr Atheroscler Rep*, 14(6), 542-555.
33. Sheikh, A., Yeasmin, F., Agarwal, S., Rahman, M., Islam, K., & Hossain, E. (2014). Protective effect of *Moringa oleifera* Lam leaves against arsenic induced toxicity in mice. *Asian Pac J Prop Biomed*, 1(8), 23-40.
34. Chattopadhyay, S., Maiiti, S., Maji, G., Deb, B., Pan, B., & Ghosh, D. (2011). Protective role of *Maringa oleifera* seed on arsenic induced hepatocellular degeneration in female albino rats. *Biol Trace Elem Res.*, 142(2), 200-212
35. Ruckmani, K., Kavimani, S., Anandan, R., & Jaykar, B. (1998). Effect of *Moringa oleifera* Lam on paracetamol - induced hepatotoxicity. *Ind J Pharmac Sci.*, 60, 33-35.
36. Gilani, A. H., Aftab, K., Suria, A., Siddiqui, S., Salem, R., & Siddiqui, B. S. (1994). Shaheen F. Pharmacological studies on hypotensive and spasmolytic activities of pure compounds from *Moringa oleifera*. *Phytoter Res*, 8(2), 87-91.
37. Dahot, M. U. (1988). Vitamin contents of flowers and seeds of *M. oleifera*. *Pak J Biochem.*, 21(4), 1-24.
38. Pal, S. K., Mukherjee, P. K., & Saha, B. P. (1995). Studies on the antiulcer activity of *Moringa oleifera* leaf extract on gastric ulcer models in rats. *Phytother Res.*, 9(6), 463-465.
39. Sreelatha, S., & Padma, P. R. (2009). Antioxidants activity and total phenolic contents of *M. oleifera* leaves in two stages of maturity plant food. *Hum Nutr.*, 64, 303-311.
40. Ofem, O. E., Ani, E. J., Archibong, A. N., & Ufford J. M. (2015). Variations in blood parameters of high salt loaded rats following administration of *Moringa oleifera* leaf extract. *Trends Med Res.*, 10(4), 97-105.
41. Archibong, A. N., Nku, C. O., & Ofem, O. E. (2017). Extract of *Moringa oleifera* attenuates hematological parameters following salt loading. *MicroMedicine*, 5 (1), 24-30.
42. CCAC. (2009). The CCAC guidelines on: The care and use of farm animals in research, teaching and testing. Canadian Council of Animal Care (CCAC), Ottawa, on [http://www.ccac.ca/en/\\_standards/guidelines/additional/fa q-farm-animals](http://www.ccac.ca/en/_standards/guidelines/additional/fa-q-farm-animals)
43. Dahiru, D. C., Onubiyi, J. A., & Umoro, H. A. (2006). Pytochemical screening and anti ulcerogenic effect of *Moringa oleifera* aqueous leaf extract. *Afr J Compl Altern Med.*, 3(3), 70-73

44. Hassarajani, S., Souzo, T., Mengi, S., & Chattopadhyaya Y. (2007). Efficacy study of the bioactive fraction (F-3) of *Acoruscalamus* in hyperlipidemia. *Indian Journal of Pharmacology*, 39, 196 – 200.
45. Obiefuna, P. C., & Obiefuna, I. P. (2001). Salt-induced hypertension in rats alters the response of isolated aortic rings to cromakalim. *West Indian Med J.*, 50, 17-21.
46. Friedewald, W. T., Levy, R. I., & Fredrickson, D. S. (1972). Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*, 18, 499-502.
47. Martirosyan, D. M., Miroshnichenko, L. A., Kulokawa, S. N., Pogojeva, A. V., & Zoloedov, V. (2007). Amaranth oil application for heart disease and hypertension. *Lipid Health Discovery*, 6(1), 1 - 4.
48. Brehm, A., Pfeiler, G., Pacini, G., Vierhapper, H., & Roden, M. (2004). Relationship between serum lipoprotein ratios and Insulin Resistance in Obesity. *Clinical Chemistry*, 50, 2316 – 2322.
49. Dabiasova, M. (2004). Atherogenic index of plasma [Log (triglyceride / HDL – Cholesterol): Theoretical and practical Implications. *Clinical Chemistry*, 50, 1113 – 1115.
50. Rice, E. C., Diplock, A. T., & Symons, M. C. (1991). Techniques I free radical research. Elsevier, Amsterdam London New York, Tokyo, 194 – 196.
51. Ohkawa, H. H., Ohishi, & Yagi, K. (1979). Assay for lipid peroxidase in animal tissues by thiobarbituric acid reaction. *Anal Biochemistry*, 95, 351 – 358.
52. Aebi, H. (1984). *Catalase in vitro Methods in Enzymology*. *Methods in Enzymology*, 105, 121–6.
53. Marklund, S. L. (1985). Pyrogallol auto-oxidation. In Handbook of methods for oxygen radical research. Ed: Greenwald R. A, Boca Raton, F. L. CRC Press, pp 43 – 247.
54. Reitman, S., & Frankel, S. (1957) Liver enzymes (AST and ALT); Reitman and Frankel calorimetric method. *Am J Uni Path.*, 28, 56.
55. Archibong, N. A., Akwari, Ad., Ofem, E. O., Irene, O. B., Samuel, U. U., & Asim, E. E. (2015). Effect of *Egeriaradiata* (Clam) Extract on Biochemical Parameters of Albino Wistar Rats. *J. Med. Sci.*, 15 (2), 87-93
56. Tiez, N. W. (1995). Clinical guide to laboratory test 3<sup>rd</sup> Edition. WB Saunders. Philadelphia PA. Pp: 528 – 519.
57. Grant, G. H. (1987). Amino acid and Protein fundamentals of clinical chemistry. Tiez N. W Editor. Third Edition. WB Saunder company Philadelphia USA, 328 – 329.
58. Drummond, K. E., & Brefere, L. M. (2014). Nutrition for food service and culinary professionals (8<sup>th</sup> edi). John Wiley and Sons.
59. Kasolo, J. (2010). Phytochemicals and uses of *M. oleifera* leave in Ugandan rural communities. *Journal of Medical plant research*, 4(9), P753.
60. Mensink, R. P., Zock, P. L., Kester, A. D., & Katan, M. B. (2003). Effect of dietary fatty acid and carbohydrate on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 control trials. *The American Journal of Clinical Nutrition*, 77 (5), 1146 – 55.
61. Hermansen, K., Dinesen, B., Hoie, L. H., Morgensten, E., & Gruenwald, J. (2003). Effect of soy and other natural products on LDL, HDL ratio and other Lipid parameters. A Literature review. *Adv. Ther.*, 20, 50 – 78.
62. Robins, H. B., Robin, S. J., Collins, D., Nelson, D. B., Elam, M. B., Schaefer, E. J., Faas, F. H., & Anderson, J. W. (2002). Diabetes plasma insulin and cardiovascular disease subgroup analysis from the department of veterans affairs. High density lipoprotein intervention trial. *Archives of internal Medicine*, 162 (22), 2597 – 604.
63. Barter, P., Gotto, A. M., Larosa, J. C., Maroni, J., Szarek, M., Grundy, S. M., Kastelein, J. J., Bittner, V., & Fruchart, J. C. (2007). HDL cholesterol, Very low level of LDL cholesterol and cardiovascular event. *The New England Journal of Medicine*, 357 (13), 1301 – 10.
64. Rahilly-Tierney, C. P., Spiro, A., Vokonas, P., & Gaziano, J. M. (2011). Relationship between high density lipoproteins cholesterol and survival to age 85 years in men. *The American Journal of Cardiology*, 107 (8), 1173 – 7.
65. Bonoy, L., Germain, S. T., Andre, H., Abene, J. H., Mahamat, M. A., & Paul, V. T. (2016). Antioxidant and antifatigue properties of aqueous extract of *M. oleifera* in rats subjected to Forced Swimming Endurance Test. *Oxidative Medicine and Cellular Longevity*, 1 – 9.
66. Moore, K., & Roberts, L. J. (1998). Measurement of lipid peroxidation". *Free Radical Research*, 28 (6), 659–71.
67. Del, R. D., Stewart, A. J., & Pellegrini, N. (2005). A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress". *Nutrition Metabolism Cardiovascular Disease*, 15 (4), 316–28.
68. Vanaporn, M., Wand, M., Michell, S. L., Sarkar-Tyson, M., Ireland, P., Goldman, S., Kewcharoenwong, C., Rinchai, D., Lertmemongkolchai, G., & Titball, R. W. (2011). Superoxide dismutase C is required for intracellular survival and virulence of *Burkholderia pseudomallei*. *Microbiology*, 157 (8), 2392–400.
69. Romao, S. (2015). Therapeutic value of oral supplementation with melon superoxide dismutase and wheat gliadin combination. *Nutrition*, 31(3), 430–6.
70. Chelikani, P., Fita, I., & Loewen, P. C. (2004). "Diversity of structures and properties among

- catalases". *Cellular and Molecular Life Sciences*, 61 (2), 192–208.
71. Li, Y., Huang, T. T., Carlson, E. J., Melov, S., Ursell, P. C., Olson, J. L., Noble, L. J., Yoshimura, M. P., Berger, C., Chan, P. H., Wallace, D. C., & Epstein, C. J. (1995). Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase. *Nature Genetics*, 11 (4), 376–81.
  72. Hamza, A. A. (2010). Ameliorative effect of *M. oleifera* Lam seed extract on liver fibrosis in rats. *Food Chemistry Toxicology*, 48, 345.
  73. Fakurazi, S., Hairuszah, I., & Nanthini, U. (2008). *Moringa oleifera* Lam prevent acetaminophen induced liver injury through the restoration of glutathione level. *Food Chemistry Toxicology*, 46, 345.
  74. Wang, C. S., Chang, T. T., Yao, W. J., Wang, S. T., & Chou, P. (2012). Impact of increasing alanine aminotransferase levels within normal range on incident diabetes". *Journal of the Formosan Medical Association*, 111 (4), 201–8.
  75. Nelson, D. L., & Cox, M. M. (2000). Lehninger, Principles of biochemistry (3<sup>rd</sup> ed) New York, Work publishing ISBN.
  76. Anderson, A. (2002). Effect of anti-inflammatory drug on the enzyme activities of the rat liver of glandular fraction which increases vascular permeability. *Journal Pharmacology*, 1, 2253 – 2254.
  77. Ezejindu, D. N., Chinweife, K. C., & Ihentuge, C. J. (2013). The effect of Moringa extract on liver enzymes of carbon tetrachloride induced hepatotoxicity in adult wister rats. *The international journal of engineering and Sciences*, 2 (7) 54 – 59.
  78. Dharmendra, S., Priya, V. A., Ved, P. & Radhey, S. G. (2014). Evaluation of antioxidant and Hepatoprotective activities of *Moringa oleifera* Lam. Leaves in carbon tetrachloride intoxicated rats. *Antioxidant*, 3, 569 – 591.
  79. Mandal, P. K., Bishayee, A. & Chatterjee, M. (1993). Stimulation of tissue repair by Mikania cordata roots extract in carbon tetrachloride induced liver injury in mice. *Phytotherapy Research*, 7, 103 – 105.
  80. Adebayo-Tayo, B. C., Onilude, A. A., & Adejoye, D. O. (2006). Bacteriological and proximate analysis of periwinkle from two different creeks in Nigeria. *World Applied Science Journal*, 1, 87-91.
  81. Burtis, C. A., Ashwood, E. R., & Bruns, T. (2006). Textbook of Clinical Chemistry and Molecular Diagnostics. 5th Edition. Elsevier, India, 621-629.
  82. Peters, T. Jr. (1996). All About Albumin: Biochemistry, Genetics and Medical Applications. Academic Press, Washington DC., USA.
  83. Bergemeyer, H. U., & Brent, E. (1974). LDU-UV assay with pyruvate and NADH. In: Method of enzymatic analysis (Ed): H. U. Bergemeyer Academic press, New York, 574 – 579.
  84. Mehta, K., Balaraman, R., Amin, A. H., Bafna, P. A., & Gulati, O. D. (2003). Effect of fruit extract of *Moringa oleifera* on Lipid profile of normal and hypercholesterolemic rabbits. *Journal ethnopharmacology*, 86(2-3), 191 – 5.
  85. Mona, S. H., Eman, M. E. & Aya, A. A. (2013). Effect of *Moringa oleifera* on serum lipid and kidney function of hyperlipidemic rats. *Journal of Applied Science Research*, 9(8), 5189 – 5198.
  86. Emmanuel, B. O., Emmanuel, G. A., Bamidele, P. F., & Martin, O. (2013). Blood glucose level and Lipid reducing activities of oral administration of aqueous leaf extract of *Moringa oleifera* on Wistar rats. *Journal of Natural Science Research*, 3 (6).
  87. Govindarajan, K., Aminu, U. K., Palanisamy, A., Norhaszalina, M. D., & Sharida, F. (2016). The modulatory effect of *Moringa oleifera* leaf extract on endogenous antioxidant system and inflammatory markers in an acetaminophen-induced nephrotoxic mice model. *Peer J.*, 4.
  88. Suaib, L., Suchita, S., Ritesh, K., Anil, K., & Debabrata, C. (2002). Experimental assessment of *M. o* leaf and fruit for its Antistress, antioxidant and scavenging potential using invitro and invivo assays. *Evidence Based Complementary and alternative Medicine*, 4(12)
  89. Nwagha, U. I., Ikekpeazu, E. J., Ejezie, F. E., Neboh, E. E., & Maduka, I. C. (2010). Atherogenic index of plasma as useful predictor of cardiovascular risk among post-menopausal women in Enugu. Nigeria. *Africa Health Science Sep.*, 10 (3), 248 – 252.
  90. Dobiasova, M., Frolalich, J., Sedova, M., Cheung, M. C., & Brown, B. G. (2011). Cholesterol esterification and atherogenic index of plasma correlate with lipoprotein size and finding on coronary angiography. *J Lipid Res.*, 52(3), 566 – 571.
  91. Dobiasova, M. (2006). AIP-atherogenic index of plasma as a significant predictor of cardiovascular risk; from research to practice.
  92. Afzal, S., Fouzia, Y., Smita, A., Mashiur, R., Khairul, I., Ekhtear, H., Shakhawoat, H., Farjana, N., Zahangir, A. S., & Khaled, H. (2014). Protective effects of *Moringa oleifera* Lam. leaves against arsenic-induced toxicity in mice. *Asian Pac J Trop Biomed.*, 4(1), 353–358.
  93. Gupta, A. K., & Misra, N. (2006). Hepatoprotective Activity of Aqueous Ethanolic Extract of Chamomile capitula in paracetamol intoxicated albino rats. *Am. J. Pharm. Toxicol.*, 1, 17 -20.