

PANCYTOPENIA-Clinico-Hematological Study in a Capital City of Telangana State -India

Dr. Majed Momin^{1*}, Dr. Abhijeet Ingle², Dr. Anamika Aluri³, Dr. K.M. Reddy G⁴, Dr. B Saroj Kumar Prusty⁵

¹Consultant pathologist, Yashoda Hospital Malakpet, Nalgonda x-roads, Hyderabad, India

²Consultant Pathologist, Yashoda Hospital Malakpet, Nalgonda x-roads, Hyderabad, India

³Consultant biochemist & HOD, Lab Malakpet, Nalgonda x-roads, Hyderabad, India

⁴General Medicine, Yashoda Hospital Malakpet, Nalgonda x-roads, Hyderabad, India

⁵Intensivist & General Medicine, Yashoda Hospital Malakpet, Nalgonda x-roads, Hyderabad, India

Original Research Article

*Corresponding author

Dr. Majed Momin

Article History

Received: 28.03.2018

Accepted: 10.04.2018

Published: 30.04.2018

DOI:

10.21276/sjmeps.2018.4.4.5



Abstract: Pancytopenia is the simultaneous presence of anaemia, leucopenia and thrombocytopenia. The etiology ranging from non-neoplastic, neoplastic to fatal leukemias. The management and prognosis depends both on the severity of pancytopenia and on the nature of underlying etiology. Thus identification of cause is the key to appropriate management. To study the clinical presentations, diagnosis and evaluation of various causes of pancytopenia by hematological parameters including bone marrow aspiration. It was prospective study. One hundred and fifty cases of Pancytopenia evaluated clinically with hematological parameters and bone marrow aspiration in a Yashoda hospital, Malakpet, Hyderabad, during period of January 2014 to February 2017. Out of 150 cases, age of patients ranged from 2 to 85 years with a mean age of 43 years. There was male preponderance. Fever and generalized weakness was common clinical presentation. The commonest physical finding was pallor and splenomegaly on systemic examination. The commonest peripheral smear was dimorphic picture & commonest bone marrow finding was hypercellular marrow with megaloblastic erythroid hyperplasia. The commonest cause for pancytopenia was Megaloblastic anemia (34%) followed by Hematolymphoid malignancy (26.6%). This study showed that clinical history, detail primary hematological investigations along with bone marrow aspiration, biopsy imprint in cytopenia patients are helpful to diagnose and to rule out the causes of pancytopenia. Megaloblastic anemia is commonest cause of pancytopenia in most indian and continent studies. Present study also shows increase in incidence of HLH (Hemophagocytic lymphohistiocytosis) and MDS are among emerging causes of pancytopenia and nutritional diet rich in vit B12 and folic acid prevents majority cause of pancytopenia in india.

Keywords: Pancytopenia, anaemia, leucopeni, thrombocytopenia, management.

INTRODUCTION

Pancytopenia is not a disease, but decreased in all three formed elements resulting from number of disease processes primarily or secondarily involving the bone marrow and symptoms attributable to anemia, leucopenia and thrombocytopenia. Pancytopenia therefore exists when hemoglobin level below 13.5g/dl in males or 11.5g/dl in females, Total leucocytes count below $4 \times 10^9/l$ and platelet counts below $150 \times 10^9/l$ [1, 2].

The etiology of pancytopenia varies in different states of Indian populations depending on the differences in age patterns, nutritional status, climate and the prevalence of infections [3]. As the severity of pancytopenia and the underlying pathology determines the management and prognosis of these patients, identifying the correct etiopathology in a given case is

crucial and helps in implementing timely and appropriate treatment [4].

The present study was done to find the common causes of pancytopenia, its clinical presentation & utility of bone marrow examination in different causes of pancytopenia in order to help the management of patients with pancytopenia attending in our hospital.

SUBJECTS AND METHODS

This is a prospective study conducted by hematopathology section, Department of Laboratory medicine, Yashoda hospital hyderabad which includes patient attending OP and admitted inpatient and diagnosed as pancytopenia during year January 2014 to February 2017. Patients of all ages and both the sexes were included. Patients with Hb <10 gm/dl, WBC <4000/cumm³, platelets <1,50, 000/cumm³ were

included. Among exclusion criterias, patient who have received or receiving chemotherapy and radiotherapy and post-transplant patient on immuno-suppressive treatment. Clinical profile includes detailed history, clinical examination and haematological parameters at presentation. Haematological profile included hemogram, peripheral blood smear morphology and bone marrow aspiration/ biopsy imprint morphology/biopsy. Blood samples of patient were obtained by routine phlebotomy procedure. 3.0 ml EDTA (K2E 5.4 mg Ethylene diamine tetra acetic acid) Vacutainer (13x75mm) anticoagulated blood was collected by needle holder and processed through Automated Hematology analyser (Beckman-Coulter LH-750). 11 Hematological parameters were considered as hemoglobin, packed cell volume, Red blood cell count, RBCs indices (MCV.MCH &MCHC), total WBCs count, Differential count, platelet counts and MPV.

A total of 150 patients who fulfilled the criteria for pancytopenia were taken up for the study. All the 150 cases were subjected to bone marrow aspiration examination after obtaining written consent from patient or guardian. Bone marrow aspiration done by Salahi's needle no [16]. From Posterior superior iliac spine or sternum and 0.3ml sample collected for morphology and 1.5 ml in EDTA and 1.5 ml in Heparin vacutainer. Leishman stain and Giemsa stains performed on aspirated sample cytosmears. Disposable Jamshidi needle (11 GX100mm) was used for bone

marrow biopsy. Bone marrow trephine biopsy was done from right posterior superior iliac spine. Whenever there is a scanty particulate marrow, blood tap or dry tap on aspiration. 1.5 to 2.0 cm long bone marrow biopsy core specimen was obtained and imprint and squash smears made and stained with leishman and giemsa stain before it fixed in fixative. Biopsy specimen was decalcified processed and sections were stained with Hematoxyline and Eosin (H&E) stain.

RESULTS

This study was carried out for 38 months in Hematopathology section, Department of laboratory medicine. A total of average 65,750 were received for Haemogram during each year. Out of these, 150 cases met inclusion criteria and bone marrow aspiration done in all cases. There were 91 males and 59 females. The mean age was 43 years with the range of 2 to 85 years. Most common affected age group was 51 to 65 years, followed by 36-50 years, 21-35 years, 02-20 years and least was more than 65 years age group (Chart 1). Male patients are predominance in almost all age groups except age group of 02-20 years were female patient found to have more pancytopenia cases.

The main chief complaints were fever, generalized weakness and loss of appetite. Among general examination pallor noted almost in all cases.




Chart-1: Age groups

Table-1: Chief complaints and Physical findings in 150 cases

| S.No | Chief complaints and Physical Findings | No. Case | % |
|------|--|----------|-------|
| 1 | Fever | 103 | 68.6 |
| 2 | Generalised Weakness | 81 | 54 |
| 3 | Shortness of breath | 37 | 24.66 |
| 4 | Headache | 15 | 10 |
| 5 | Giddiness | 11 | 7.33 |
| 6 | Blurring of Vision | 03 | 2 |
| 7 | Loss of weight | 12 | 8 |
| 8 | Loss of Apetite | 41 | 27.33 |
| 9 | Body pain | 21 | 14 |
| 10 | Cough | 25 | 16.66 |
| 11 | Bleeding Manifestation | 12 | 8.0 |
| 12 | Vomittings | 18 | 5.33 |
| 13 | Loose Motion | 07 | 4.66 |
| 14 | Pallor | 150 | 100 |
| 15 | Jaundice | 16 | 10.66 |
| 16 | Lymphadenopathy | 07 | 4.66 |
| 17 | Oedema over feet | 08 | 5.33 |
| 18 | Hepatomegaly | 23 | 10.66 |
| 19 | Splenomegaly | 55 | 36.66 |

Splenomegaly was more commonly seen than hepatomegaly (Table 1).

Hematological parameters as automated hematology analyzer sub grouped in to mild, moderate and severe form are shown in (Table: 2).

Table-2: Grading of formed element

| Parameters | Range | Cases | % |
|---------------------------------|----------------------|-------|-------|
| Hemoglobin | | | |
| Severe | 21 to 5.5 gm% | 43 | 28.7% |
| Moderate | 5.6 to 8.0 gm% | 65 | 43.3% |
| Mild | 8.1 to 10 gm% | 42 | 28% |
| White blood cells (WBCs) | | | |
| Severe | 100 to 1000/cumm | 15 | 10% |
| Moderate | 1100 to 2500/cumm | 68 | 45.3% |
| Mild | 2600to 4000/cumm | 67 | 44.7% |
| Platelets | | | |
| Severe | Less than30000/cumm | 44 | 29.3% |
| Moderate | 31000 to 80000/cumm | 68 | 45.3% |
| Mild | 81000 to 140000/cumm | 38 | 25.4% |

The most common peripheral blood RBCs picture was Dimorphic picture (28%). Hyper segmented neutrophils seen in 95% cases in megaloblastic anemia and 05% in mixed deficiency cases. Leucopenia and thrombocytopenia were seen in all cases.

Out of 150 cases of pancytopenia .the most common cause was Megaloblastic anemia (34%) (Table: 3).




Chart 2
 Dimorphic RBCs28%
 Normocytic Normochromic RBCs26%
 Macrocytic RBCs24%
 Normocytic Hypochromic RBCs12%
 Microcytic Hypochromic RBCs10%

Table-3: Distribution of Different etiology of Pancytopenia among 150 cases

| Sr.no | Etiology | Number of cases | Percentage % | M | F | M:F |
|-------|---|-----------------|--------------|----|----|-------|
| 1 | Megaloblastic anemia | 51 | 34 | 34 | 17 | 2:1 |
| 2 | Acute leukemia | 21 | 14 | 09 | 12 | 0.7:1 |
| 3 | Lymphoproliferative disorder | 12 | 08 | 07 | 04 | 1.7:1 |
| 4 | Myelodysplastic syndrome (MDS) | 07 | 4.66 | 05 | 02 | 2.5:1 |
| 5 | Aplastic anemia | 17 | 11.33 | 14 | 03 | 4.6:1 |
| 6 | Hypersplenism | 13 | 8.66 | 04 | 09 | 0.4:1 |
| 7 | Plasma cell dyscrasia | 05 | 3.33 | 03 | 02 | 1.5:1 |
| 8 | Hemophagocytic lymphohistiocytosis(HLH) | 04 | 2.66 | 03 | 01 | 3:1 |
| 9 | Disseminated Tuberculosis | 03 | 2 | 01 | 02 | 0.5:1 |
| 10 | Disseminated carcinomas | 02 | 1.33 | 01 | 01 | 1:1 |
| 11 | Mixed deficiency anemia | 06 | 4.0 | 04 | 02 | 2:1 |
| 12 | Hemolytic anemias | 03 | 2 | 03 | 00 | 3:0 |
| 13 | Sepsis | 04 | 2.66 | 01 | 03 | 0.3:1 |
| 14 | SLE | 01 | 0.66 | 01 | 00 | 1:0 |
| 15 | Drug induced | 01 | 0.66 | 01 | 00 | 1:0 |


Megaloblastic anemia found in 51 cases with male predominance. Hemogram RBCs indices in megaloblastic anemia show average MCV 105fl (76-94fl) and MCH was 35pg (27-32 pg). Vitamin B12 deficiency alone more pronounced 78%, folic acid deficiency 04% and 18% cases in mixed deficiency. Hyper segmented neutrophils and macrovalocytes (Figure 1a) are predominant peripheral smear picture. Bone marrow aspirate mostly hyper cellular with increased erythroid: myeloid ratio. Megaloblasts (Figure 1b; red arrow) are large with sieve like nuclear chromatin, bluish cytoplasm. Giant metamyelocytes and stab forms (Figure 1b; blue arrow) commonly associated finding in aspirate smears.

Among Hematolymphoid malignancy, acute promyelocytic leukemia and acute monoblastic leukemia (figure 2) commonly encountered. Diluted marrow seen in acute lymphoblastic leukemia and imprint biopsy smears helps in early morphological diagnosis of leukemia. Erythroid and megakaryocytic series reduced in majority of cases of hematolymphoid malignancy.

Aplastic anemia commonly encountered in male and Bone marrow biopsy performed in all cases.

Reactive lymphoplasmacytosis seen with increased fat spaces (figure3). Hypersplenism surprisingly found more in female. Significant Rouleaux formation and Erythrocyte sedimentation rate with A: G reversal seen in almost all cases (figure 4). Hem phagocytic lymphohistiocytosis (HLH) seen two paediatric age group of 4 and 13years, other two cases at 60 year and 39 year male (figure6). Hepato-splenomegaly, high ferritin, hypertriglyceridemia and increased lactate dehydrogenase seen all cases. Bone marrow granulomas (figure5) of tuberculus origin seen in two cases. Cervical tuberculosis seen in both the cases. Disseminated carcinoma seen in one female patient from carcinoma breast and other male patient had prostatic adenocarcinoma (figure7) deposit. Serum PSA was very high (1650 ng/ml).

We encountered 4 cases of sepsis with female predominance. Neutrophils toxic granules and cytoplasmic vacillations seen in these cases. Plasma procalcitonin level is more than 75 ng/ml on an average. Hemolytic anemia cases show reticulocytosis and positive coombs test. One case diagnosed as Evans syndrome.

**Fig-1: a, bMegaloblastic anemia (PS&BM images)**




Fig-2: a, b acute leukemia (ALL &AML)




Fig-3: Aplastic anemia (BM aspiration & biopsy)




Fig-4: Multiple myeloma (PS & BM)




Fig-5: Bone marrow granulomas




Fig-6: Hemophagocytic lymphohistiocytosis




Fig-7: Disseminated carcinomas

Figure 1a: Peripheral smear (Leishman stain 40X) shows Macrovalocytes (blue arrow) & hypersegmented neutrophils (Red Arrow).
Figure 1b: Bone marrow aspirate smear (Giemsa stain 100X) Megaloblast with sieve like chromatin (Red arrow) & Giant stab forms (blue arrow).
Figure 2a: Peripheral smear (Leishman stain 40X) lymphoblast in peripheal smear,2b: lymphoblast in Bipsy imprint,2c: myeloblast in peripheral smear,2d: Atypical promyelocyte with auer rod.
Figure 3a: Bone marrow aspirate smear (Giemsa stain 100X) Increased fat with reactive plasmacytosis; 3b: Bone marrow biopsy (H&E Stain, 40X) increased fat spaces.
Figure 4a: Peripheral smear (Leishman stain 40X) show significant roulaux formation.
Figure 4b: Bone marrow aspirate smear (Giemsa stain 100X) Many plasma cells with eccentric nuclei.
Figure 5a: (Leishman stain 40X) Bone marrow aspirate granulomas with epitheloid cells. 5b; Peripheral smear (Leishman stain 40X) show leucoerythroblastic picture. 5c; Bone marrow biopsy (H&E Stain, 40X) bone marrow granulomas.
Figure 6a: Bone marrow aspirate smear (Giemsa stain 100X) Hemophagocytes .6b; Bone marrow biopsy (H&E Stain, 40X) hemophagocytosis.
Figure 7a: Bone marrow aspirate smear (Leishman stain 40X) tumour cells in loose cohesive clusters .7bBone marrow biopsy (H&E Stain, 40X) adenocarcinoma cells from prostate.

DISCUSSION

During period of three years and two months, 150 cases of pancytopenia studied. Different etiology of pancytopenia studied and compared with the national and little international published literature.

The mean age was 43 years in a patients ranged from 2 to 85 years ago with male predominance (60%), female (40%) and male to female ratio (M: F) of 1.5:1. Age and sex distribution was compared with other studies.

Table-4: Age, sex distribution compared to other studies done on pancytopenia

| sr.no. | Authours | Country | Year | Number of case | Age ranges | M:F Ratio |
|--------|-------------------------------|----------|------|----------------|------------|-----------|
| 1 | Khunger <i>et al.</i> [4] | India | 2002 | 200 | 02-70 | 1.2:1 |
| 2 | Ashraf <i>et al.</i> [5] | Pakistan | 2010 | 150 | 15-60 | 1.1:1 |
| 3 | Gayathri BN <i>et al.</i> [6] | India | 2011 | 104 | 02-80 | 1.2:1 |
| 4 | Vandana <i>et al.</i> | India | 2012 | 80 | 01-79 | 1:1.2 |
| 5 | Sweta <i>et al.</i> [7] | India | 2014 | 100 | 05-80 | 1.5:1 |
| 6 | Present study | India | 2017 | 150 | 02-85 | 1.5:1 |

Fever was the most common presenting complaint in our study (68.6%) followed by Generalized weakness (54%). The most common physical finding was pallor (100%), followed by splenomegaly (36%). Peripheral blood smear picture RBCs morphology Macrocytic and dimorphic together constitute 50%, followed by Normocytic normochromic RBCs (28%), Normocytic hypochromic RBCs (12%) & microcytic hypochromic (10%). Leucoerythroblastic picture seen (12.6%) with dimorphic picture and commonly associated with megaloblastic anemia and infiltrative neoplastic lesion.

WBCs differential count lymphocyte predominance noted in 65% of hypoplastic anemia in our study compared to 50% in Tilak *et al.* [11], 51.3% in Gaythri & Rao *et al.* [6] and 85% in Khunger JM *et al.* [4] study. Hypersegmented neutrophils seen in 90% of cases which was highest compared to 84.9% in Tilak V *et al.* [11] Study, 51.3% in Gayathri *et al.* [6]. Platelet anisocytosis, large and giant platelets more pronounced in Myelodysplastic syndrome followed by megaloblastic anemia.

The commonest cause of pancytopenia reported in our study was megaloblastic anemia, similar to other studies conducted in asian countries. While commonest cause of pancytopenia in various studies throughout world has been aplastic anemia. In our study aplastic anemia was fifth common cause of pancytopenia.

All indian studies suggest more incidence of megaloblastic anemia among all etiology of pancytopenia. Incidence of megaloblastic anemia in our study was 34%, incidence of 74% in Gayathri & Rao *et al.* 72% was reported by Khunger JM *et al.* [4], 68% by Tilak *et al.* [11]. All these indian studies highlights higher prevalence of nutritional anemia in indians. Although Histogram RBCs Indices (MCV, MCH) & Peripheral smear examinations are more informative. Bone marrow aspiration performed to rule out possibility of marrow dysplasia. In our study serum Vitamin B12 and folic acid estimation done in those cases who are not treated with injectable or oral supplement before admission. Vitamin B12 deficiency found in 86.1% & folic acid 14% cases.

Table-5: Major causes of pancytopenia compare to other studies

| Studies | Country | Year of study | Number of cases | Commonest cause | Second commonest cause |
|---------------------------------|-----------------|---------------|-----------------|------------------------------------|-----------------------------------|
| Keisu and Ost [8] | Israel & Europe | 1990 | 100 | Neoplastic disease radiation (32%) | Hypoplastic anemia (19%) |
| Hossain <i>et al.</i> [9] | Bangladesh | 1992 | 50 | Hypoplastic anemia | Chronic malaria and kalazar |
| Jha <i>et al.</i> [10] | Nepal | 2005 | 148 | Hypoplastic anemia (29%) | Megaloblastic anemia (23%) |
| Ashraf S <i>et al.</i> [5] | Pakistan | 2010 | 150 | Hypersplenism (68%) | Megaloblastic anemia (25.4%) |
| Tilak & Jain <i>et al.</i> [11] | India | 1999 | 77 | Megaloblastic anemia (68%) | Hypoplastic anemia (7.7%) |
| Kumar <i>et al.</i> [3] | India | 2001 | 166 | Aplastic anemia (29.5%) | Megaloblastic anemia (22.3%) |
| Gayathri BN <i>et al.</i> [6] | India | 2011 | 104 | Megaloblastic anemia (74%) | Aplastic anemia (18.3%) |
| Vandana R <i>et al.</i> | India | 2012 | 80 | Megaloblastic anemia (41.2%) | Dimorphic anemia (8.7%) |
| Sweta <i>et al.</i> [7] | India | 2014 | 100 | Megaloblastic anemia (66%) | Aplastic anemia (18%) |
| Vaddati T <i>et al.</i> [12] | India | 2015 | 75 | Megaloblastic anemia (56%) | Aplastic anemia (13.3%) |
| Present study | India | 2017 | 150 | Megaloblastic anemia (34%) | Hemato-lymphod malignancy (26.6%) |

We encountered 26.6 % of hematolymphoid malignancy presenting as pancytopenia. Out of which 14% was acute leukemia, 08% of lymphoproliferative disorder and 4.6% of myelodysplastic syndrome. Incidence reported by Khunger JM *et al.* [4] (5%), 12 % Reported by Kumar *et al.* [3] and 3.85% by Gayathri and rao *et al.* [6]. Dry tap diluted marrow encountered in 20% cases. Bone marrow biopsy imprint smears play crucial role in early diagnosis especially in diluted and dry tap marrow. The diagnosis further confirmed on flow cytometry immune phenotyping and molecular study.

The incidence of Hypoplastic marrow in our study was 11.3% which very close to studies done by khunger *et al.* [4] (14%). While higher incidence reported by Kumar *et al.* (29.5%) [3]. Bone marrow biopsy done almost all cases of hypoplastic marrow. The cause of hypoplastic was idiopathic in 90% of cases. Methotrexate induced hypoplastic marrow seen in three cases.

Plasma cell dyscrasia constituting 3.33% compared to 4% reported by Khodke *et al.* 1.3% by Tilak *et al.* [11] and Khunger *et al.* [4] who have reported an incidence of 1% in their studies. All cases show significant rouleaux formation on peripheral blood film and average ESR of 100 mm at the end of one hr by Westergrens method. Biochemical serum protein show A: G reversal seen in almost all cases with normal alkaline phosphatase levels.

We encountered four cases of Hemophagocytic lymphohistiocytosis (HLH). Two cases related to tuberculosis and other two was idiopathic. Major and minor criteria were considered for final diagnosis of HLH and 50 % Mortality seen in our HLH.

Disseminated Tuberculosis and Disseminated carcinomas share incidence 2.0% and 1.33% respectively. Leucoerythroblastic picture seen in one case each. Prostate carcinoma infiltration observed in one case and breast carcinoma in other case of disseminated carcinomas and confirmed with bone marrow biopsy immunohistochemistry. Disseminated carcinoma was not found in other studies done by Khunger *et al.* [4], Khodke *et al.* Tilak *et al.* [11] and Gayathri & Rao *et al.* [6] Kumar *et al.* [3] reported 2 cases of marrow metastasis in his studies. Hemolytic anemia and sepsis share around 2% incidence while SLE was rare cause of pancytopenia in our studies.

The cause of pancytopenia was treatable in almost all cases of megaloblastic anemia. Mortality found in case of hematolymphoid malignancy, HLH and severe aplastic anemia, due to overwhelming infections.

CONCLUSION

Pancytopenia is not a disease entity. Prolonged fever, generalized weakness, prolonged anemia and tendency to bleed could be suspected for pancytopenia on clinical ground. Megaloblastic anemia was the most common cause of pancytopenia, correlating with other studies done in India. Incidence of HLH and MDS emerging etiology compare to other studies. Haemogram with RBCs indices, Peripheral blood smear morphology, Bone marrow aspiration cytology and biopsy imprint smears morphology would be helpful in early identification and early intervention of patient with pancytopenia which helps in understanding disease process and early planning for further investigations for accurate diagnosis and management, to enhance survival rate in case of pancytopenia.

REFERENCES

1. Firkin, F., Chesterman, C., Rush, B., & Pennigton, D. (2008). *De Gruchy's Clinical haematology in medical Practice*. John Wiley & Sons.
2. Guinan, E. C. (2004). Acquired and inherited aplastic anemia syndromes. *Wintrobe's Clinical Hematology*, 1396-1419.
3. Kumar, R., Kalra, S. P., Kumar, H., Anand, A. C., & Madan, H. (2001). Pancytopenia--a six year study. *The Journal of the Association of Physicians of India*, 49, 1078-1081.
4. Gayathri, B. N., & Rao, K. S. (2011). Pancytopenia: a clinico hematological study. *Journal of laboratory physicians*, 3(1), 15.
5. Ashraf, S., & Naeem, S. (2010). Frequency of hypersplenism in chronic liver disease patients presenting with pancytopenia. *Annals of King Edward Medical University*, 16(1 SI).
6. Jain, A., & Naniwadekar, M. (2013). An etiological reappraisal of pancytopenia--largest series reported to date from a single tertiary care teaching hospital. *BMC Blood Disorders*, 13(1), 10.
7. Barik, S., Chandoke, R. K., & Verma, A. K. (2014). A prospective clinico-hematological study in 100 cases of pancytopenia in capital city of India. *Journal of applied hematology*, 5(2), 45.
8. Keisu, M., & Öst, Å. (1990). Diagnoses in patients with severe pancytopenia suspected of having aplastic anemia. *European journal of haematology*, 45(1), 11-14.
9. Hossain, M. A., Akond, A. K., Chowdhary, M. K., Sikder, A. M., & Rashid, M. A. (1992). Pancytopenia-A study of 50 cases. *Bangladesh Journal of pathology*, 1, 9-12.
10. Jha, A., Sayami, G., Adhikari, R. C., Panta, A. D., & Jha, R. (2008). Bone marrow examination in cases of pancytopenia.
11. Tilak, V., & Jain, R. (1999). Pancytopenia--a clinico-hematologic analysis of 77 cases. *Indian journal of pathology & microbiology*, 42(4), 399-404.
12. Tejeswini, V., Premalatha, P., Renuka, I. V., Krishna, R., Krichnamacharyulu, P. A. V., &

Vahini, G. (2015). Clinicohaematological profile of pancytopenia-A South indian tertiary hospital experience. *Indian Journal of Pathology and Oncology*, 2(3), 165-169.