

A Study on the Prevalence of *Cysticercus cellulose* in Pigs of Son La Province, Vietnam

Nguyen Thi Kim Lan*, Do Thi Lan Phuong, Phan Thi Hong Phuc, Pham Dieu Thuy, Dao Van Cuong
Faculty of Animal Husbandry and Veterinary Medicine, Thai Nguyen University of Agriculture and Forestry, Thai Nguyen city, Thai Nguyen prefecture, Vietnam

Original Research Article

***Corresponding author**
Nguyen Thi Kim Lan

Article History

Received: 01.04.2018
Accepted: 13.04.2018
Published: 30.05.2018

DOI:

10.21276/sjmeps.2018.4.5.17



Abstract: Human taeniasis was found from more than 50 provinces in Vietnam, and in which the highest prevalences have always been observed in the highland and mountainous areas. In addition, *Cysticercus cellulose* infection of pigs is considered as one of the most risk factor that contribute to taeniasis in human. Therefore, our study investigated both of the prevalence of *Cysticercus cellulose* in pigs and *Taenia solium* in human in the 3 communes of Son La, a mountainous province, in order to identify if there is such kind of correlation between the infected rates of pigs and human. Our results show that the prevalences of *Cysticercus cellulose* in pigs and *Taenia solium* in human in the 3 communes were high when comparing to other areas of Vietnam, however, they are within the range of the disease infected rates in Vietnamese mountainous areas. In addition, we observed that infection of *Cysticercus cellulose* in pigs was closely associated by infection of *Taenia solium* in human, and also vice versa. In addition, because although cysticerci of *Taenia solium* occur primarily in pork, they also occur in humans, so humans can have both taeniasis and cysticercosis (including neurocysticercosis), which are dangerous and health-damaged in many cases. Therefore we suggest that in order to eliminate this meat-borne parasitoses, the effective prevention and treatment are required to be applied simultaneously on both pigs and humans of the areas.

Keywords: *Cysticercus cellulose*, *Taenia solium*, Vietnam, mountainous area.

INTRODUCTION

Taeniasis and cysticercosis are endemic in Africa, South and Central America, Brazil, Mexico, China, India, Myanmar, Malaysia, Korea, Indonesia, Philippines, and Southeast Asia, including Vietnam [1-3]. It has been estimated that hundreds of millions of persons worldwide are infected with *Taenia solium*, the most serious tapeworm species to humans [4, 1, 5]. Human taeniasis was found from more than 50 provinces in Vietnam [3]. In addition, the highest prevalence was in the highland and the mountainous areas [6,7]. In our study, we investigated the prevalence of *Cysticercus cellulose* in pigs and *Taenia solium* in human in the 3 districts of Son La province, a mountainous area, in order to identify if there is a

relationship between the infected rates on pigs and human.

MATERIALS AND METHODS

The prevalences of *Cysticercus cellulose* were identified in 1040 pigs in Bac Yen, Muong La and Mai Son communes, by the exploratory surgery method. The prevalences of *Taenia solium* were diagnosed in 750 people of the same communes. We identified the significance of correlation between the infected rates by evaluating the Pearson correlation coefficient value.

RESULTS AND DISCUSSION

The prevalences of *Cysticercus cellulose* in pigs of the 3 communes of Son La province are shown in Table-1 and Photo 1-5.

Table-1: The prevalence of *Cysticercus cellulosae* in pigs of the 3 communes of Son La province

Commune	Number of diagnostic pigs	Number of infected pigs	Prevalence (%)
Bac Yen	351	11	3.13
Muong La	337	9	2.67
Mai Son	352	7	1.99
Total	1,040	27	2.59



Photo-1: *Cysticercus cellulosae* in pig muscle




Photo-2: *Cysticercus cellulosae* in pig brain




Photo-3: *Cysticercus cellulosae* in pig heart



Photo-4: *Cysticercus cellulosae* in pig tongue



Photo-5: Isolated *Cysticercus cellulosae* from pig organs

From Table-1, we observe that the prevalence of *Cysticercus cellulosae* in pigs of the 3 communes are relatively high, from 1.99 to 3.13 %. The prevalences are much higher than those identified in the northern and the southern slaughter houses in Vietnam, from 0.03 to 0.9 % [7, 8]. It can be explained by the fact that Son La is a mountainous area with the undeveloped economic situation, in which pigs are usually let

wandering freely outside in daytime, and therefore they are easily to get access to the parasitic contaminated sources.

We also identified the prevalences of *Taenia solium* in human of the 3 communes of Son La province, and the results are shown in Table-2 and Photo 6.

Table-2: The prevalences of *Taenia solium* in human of the 3 communes of Son La province

Commune	Number of diagnostic humans	Number of infected humans	Prevalence(%)
Bac Yen	250	10	4.00
Muong La	250	8	3.20
Mai Son	250	5	2.00
Total	750	23	3.07



Photo-6: *Taenia solium* infected in human

From Table-2, we observe that the prevalence of *Taenia solium* in human of the 3 communes are relatively high, from 2.00 to 4.00 %. However, the prevalences are within the normal infected range of the mountainous areas in Vietnam recorded in several

previous research, which reporting the rates of from 2.0 to 10.4% [6, 7]. Unhygienic toilet systems in mountainous areas, such as shown in photo 7, is usually considered as an important reason that contributing to the high prevalence of *Taenia solium* in human.



Photo-7: A typical human toilet in Son La province

In order to identify the relationship, we computed the linear regression and calculated the Pearson correlation coefficient value between the

infected rates of *Cysticercus cellulose* in pigs and of *Taenia solium* in human of the 3 communes of Son La province, and the results shown in Figure-1.




Fig-1: The computed linear regression between the infected rates of *Cysticercus cellulose* in pigs and of *Taenia solium* in human of the 3 communes of Son La province

From Figure-1, we observe that the Pearson correlation coefficient value (R) was significant and close to 1 (R = 0.929), suggesting that infection of *Cysticercus cellulose* in pigs is highly associated by infection of *Taenia solium* in human and vice versus. Although cysticerci of *Taenia solium* occur primarily in pork, they also occur in humans. Therefore, humans can have taeniasis and cysticercosis (including neurocysticercosis), which is dangerous and health-damaged in many cases [9, 2, 3]. In order to eliminate this meat-borne parasitoses, the prevention and treatment methods are required to simultaneously applied in both pigs and people of the areas.

ACKNOWLEDGMENTS

We (corresponding author: Nguyen Thi Kim Lan and all other members of the group) express sincere thanks to Vietnam Ministry of Education & Training for funding this research, under the grant number B2017-TNA-34.

CONFLICT OF INTEREST

We have no conflict of interest related to this study.

REFERENCES

1. Ito, A., Nakao, M., & Wandra, T. (2003). Human taeniasis and cysticercosis in Asia. *The Lancet*, 362(9399), 1918-1920.
2. Flisser, A., Geerts, S., Kyvsgaard, N. C., McManus, D. P., Nash, T. E., & Pawlowski, Z. (2005). *FAO/WHO/OIE Guidelines for the Surveillance, Prevention and Control of Taeniosis/cysticercosis* (No. V460 MURw). K. D. Murrell, & P. Dorny (Eds.). Paris: World Organization for Animal Health.
3. Van De, N., Le, T. H., Lien, P. T. H., & Eom, K. S. (2014). Current status of taeniasis and cysticercosis in Vietnam. *The Korean journal of parasitology*, 52(2), 125.
4. Eom, K. S., & Rim, H. J. (1993). Morphologic descriptions of *Taenia asiatica* sp. n. *Korean J Parasitol*, 31(31), 1-6.
5. Nakao, M., Okamoto, M., Sako, Y., Yamasaki, H., Nakaya, K., & Ito, A. (2002). A phylogenetic hypothesis for the distribution of two genotypes of the pig tapeworm *Taenia solium* worldwide. *Parasitology*, 124(6), 657-662.
6. Van De, N., Le, T. H., Lien, P. T. H., & Eom, K. S. (2014). Current status of taeniasis and cysticercosis in Vietnam. *The Korean journal of parasitology*, 52(2), 125.
7. De, N. V., & Le, T. H. (2010). *Taenia/cysticercosis and molecular application* (textbook). Hanoi, Vietnam.
8. Trieu, H. S. (2012). *Study on genotype of pathogen, clinical, sub-clinical symptoms, treatment efficacy for taeniasis and cysticercosis patients in National Institute of Malariology, Parasitology and*

Entomology 2007-2010 (Doctoral dissertation, PhD thesis. 2012 (in Vietnamese)).

9. Fan, P. C., & Chung, W. C. (1998). *Taenia saginata asiatica: Epidemiology, infection, immunological and molecular studies. Journal of microbiology, immunology, and infection= Wei mian yu gan ran za zhi*, 31(2), 84-89.